

---

**ISSN 2349-7750****INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**

---

Available online at: <http://www.iajps.com>

---

**Review Article****BIOLOGICAL AND PHARMACEUTICAL BUFFER SYSTEMS****M. Hareesh Reddy\*, K. Hariprasad Rao, V. Muralidharan, N. Muralikrishna**

Shadan College of Pharmacy, Peerancheru, Hyderabad, Telangana.

**Abstract:**

Generally, buffers are used in the pharmaceutical products for two purposes, viz., to adjust the PH of Product for maximum stability and to maintain the PH within the optimum physiological PH range. Pharmaceutical solutions generally have a low buffer capacity in order to prevent overwhelming the bodies' own buffer systems and significantly changing the PH of the body fluids. Buffers have concentrations in the range of 0.05 to 0.5M and buffer capacities in the range of 0.01 to 0.1, which are usually sufficient for pharmaceutical solutions. Most pharmaceutical buffers are composed of ingredients that are found in the body (eg. acetate, phosphate, citrate and borate). The ingredients are selected from the available ones considering their sterility, stability, cost, toxicity, etc.

**Key Words:** *Buffers, Optimization, Citric acid, Lacrimal fluids.*

**Corresponding Author:**

**M. Hareesh Reddy,**  
Dept. of Pharmaceutics,  
Shadan College of Pharmacy,  
Peerancheru, Hyderabad

QR code



---

Please cite this article in press as Hareesh Reddy *et al.* **Biological and Pharmaceutical Buffer Systems, Indo American J of Pharm Sci, 2015;2(2):563-566.**

## BUFFERS IN PHARMACEUTICAL SYSTEMS

### Solid Dosage Forms:

Buffers have been widely used in solid dosage forms such as tablets, capsules and powders for controlling the pH of the environment around the solid particles. This has practical application for the drugs that have dissolution rate limited absorption from unbuffered solutions [1-6]. One of the special applications of buffers is to reduce the gastric irritation caused by the acidic drugs. For example, sodium bicarbonate, magnesium carbonate and sodium citrate antacids are used for the purpose of reducing toxicity [7-10].

### Semisolid Dosage Forms:

Semisolid preparations such as creams and ointments undergo pH changes upon storage for long storage for a long time, resulting in its reduced stability. Hence; buffers such as citric acid and sodium citrate or phosphoric acid/sodium phosphate are included in these preparations so as to maintain their stability.

### Parenteral Products:

Use of buffers is common in the parenteral products. Since the pH of blood is 7.4, these products are required to be adjusted to this pH. Change in pH to higher side (more than 10) may cause tissue necrosis while on lower side (below 3) it may cause pain at the site of action. Commonly used buffers include citrate, glutamate, phthalate and acetate. The pH optimization is generally carried out to have better solubility, stability and reduced irritancy of the product.

### Ophthalmic Products:

The purpose of buffering some ophthalmic solution is to prevent an increase in pH can affect both the solubility and the stability of the drug. The decision whether or not buffering agents should be added in preparing an ophthalmic solution must be based on several considerations. Normal tears have pH of about 7.4 and possess some buffer capacity [11-14].

## BUFFERS IN BIOLOGICAL SYSTEMS

Buffers for biological systems should satisfy the following requirements

- A pKa value between 6.0 to 8.0
- High water solubility and minimal organic solvent solubility

- Exclusion by cellular membranes
- Minimal interaction between buffer and critical reaction compounds
- Should not absorb light in the visible and UV spectrum
- Stable and resistant to enzymatic degradation
- Minimal changes on dissociation from changes in concentration and temperature
- Easy to manufacture and purify
- Cost effective
- Ionic strength
- Permeability

Buffers are important to biological systems because they enable the organism to maintain homeostasis with the little physiological cost

**Blood:** Blood consists of primary and secondary buffer systems contributing the pH 7.4. When pH of the blood goes below 7.0 or above 7.8, life is in danger. The pH of the blood in diabetic coma is reported to drop as low as 6.8.

Primary buffers present in plasma are – Carbonic acid-bicarbonate system and acid alkali salts of phosphoric acid systems

Secondary buffers that are present in erythrocytes are – hemoglobin/oxyhemoglobin system and acid salts of phosphoric acid system

**Lacrimal Fluids:** Lacrimal fluids (tears) have been found to have a great degree of buffer capacity, allowing dilution of 1:15 with neutral distilled water. The pH of tears is about 7.4, with a range of 7.0 to 8.0. Normally, pure conjunctival fluid is more acidic than the tear fluid commonly employed in pharmacy. The pH increases rapidly when the sample is removed for analysis because of loss of carbon dioxide from the tear fluid

## Three Buffer Systems in Body Fluid

### Protein Buffer Systems:

Proteins are the most important and widely operating buffers in the body fluid. The protein buffer system is an integral component of the body's pH controlling mechanism. Protein buffers are either intracellular or extracellular [15-19]. Their functionality is mainly intracellular focused and include haemoglobin (Hb). Hb is the protein that functions to transport oxygen within the body. Plasma proteins function as buffers but their amount is small in comparison with the

intracellular protein buffers. Protein buffers include basic group, and acidic protein buffer groups, that act as hydrogen ion depleters or donors to maintain the pH level at 7.4. The most well-known protein buffers include 0.1 M NaH<sub>2</sub>PO<sub>4</sub>, pH 6.2 (Activation buffer), PBS, pH 7.4 (Alternate Coupling Buffer) and the PBS, 1 percent BSA, pH 7.4 (Assay Buffer).

#### Phosphate Buffer System [20-22].

The phosphate buffer system is comprised of two ions: hydrogen phosphate ions and dihydrogen phosphate ions. The pH level of the blood drops below 7.4 when the H<sup>+</sup> ions in the bloodstream increase. Hydrogen phosphate ions accept all additional H<sup>+</sup> ions to reestablish the equilibrium between the hydroxide and hydrogen ions in the blood. When the pH level of the blood increases above 7.4, the dihydrogenate phosphate ions release additional hydrogen ions to reinstate the pH level of the blood to its optimal 7.4.


#### Bicarbonate Buffer System [23-25]

The bicarbonate buffer system functions to maintain the pH level in the blood of mammals. It also plays a major role in the formation of acid in the stomach, and to neutralize the pH of chyme that enters the small intestine from the stomach. The bicarbonate

buffer system manages acid/base imbalances and effectively manages the release of excess carbon dioxide as a bi-product of cellular respiration.

#### BUFFER CAPACITY [26-29]

Buffering capacity is defined as the number of moles of strong base or acid needed to change the pH of a liter of buffer solution by one unit. A general buffer capacity estimate is 40 percent of the total sum of the molarities of the conjugate base and acid.



**Table1: Standard Buffers with their pH Range and Quantities of Ingredients**

Buffer	pH	Method
Hydrochloric acid Buffer	1.2 - 2.2	50 ml of 0.2M kcl and a sufficient amount of 0.2 N Hcl. Final volumes is made by adding water to make 200ml solution.
Acid phthalate buffer	2.2 - 4.0	50 ml potassium hydrogen phthalate and sufficient volume of 0.2N Hcl. Final volume is made by adding water to make 200 ml solution
Neutralized phosphate buffer	4.2 - 5.8	50 ml potassium hydrogen phthalate and sufficient volume of 0.2N NaOH. Final volume is made by adding water to make 200 ml solution
Phosphate buffer	5.8 - 8.0	50 ml potassium dihydrogen phosphate and sufficient volume of 0.2N Hcl. Final volume is made by adding water to make 200 ml solution
Alkaline borate buffer	8.0 - 10.0	50 ml boric acid-potassium chloride and sufficient volume of 0.2N NaOH. Final volume is made by adding water to make 200 ml solution

#### REFERENCES

- Steptoe PC, Edwards RG, Purdy JM. Human blastocysts grown in culture. *Nature*. 1971;229(5280):132–3.
- Brinster RL. Studies on the Development of Mouse Embryos in Vitro. I. The Effect of Osmolarity and Hydrogen Ion Concentration. *J Exp Zool*. 1965;158:49–57.
- Hershlag A, Feng H. The effect of CO<sub>2</sub> concentration and pH on the in vitro development of mouse embryos. *Fertil Magazine*. 2001;4:21–2.
- Carney EW, Bavister BD. Regulation of hamster

- embryo development in vitro by carbon dioxide. *Biol Reprod.* 1987;36(5):1155–63.
5. Hentemann M, Mousavi K, Bertheussen K. Differential pH in embryo culture. *Fertil Steril.* 2011;95(4):1291–1294.
  6. Emmens CW. The motility and viability of rabbit spermatozoa at different hydrogen-ion concentrations. *J Physiol.* 1947;106(4):471–481.
  7. Pholpramool C, Chaturapanich G. Effect of sodium and potassium concentrations and pH on the maintenance of motility of rabbit and rat epididymal spermatozoa. *J Reprod Fertil.* 1979;57(1):245–251.
  8. Bagger PV, Byskov AG, Christiansen MD. Maturation of mouse oocytes in vitro is influenced by alkalinization during their isolation. *J Reprod Fertil.* 1987;80(1):251–255.
  9. Downs SM, Mastropolo AM. Culture conditions affect meiotic regulation in cumulus cell-enclosed mouse oocytes. *Mol Reprod Dev.* 1997;46(4):551–566.
  10. Leclerc C, Becker D, Buehr M, Warner A. Low intracellular pH is involved in the early embryonic death of DDK mouse eggs fertilized by alien sperm. *Dev Dyn.* 1994;200(3):257–267.
  11. Zhao Y, Baltz JM. Bicarbonate/chloride exchange and intracellular pH throughout preimplantation mouse embryo development. *Am J Physiol.* 1996;271(5 Pt 1):C1512–1520.
  12. Lane M, Baltz JM, Bavister BD. Na<sup>+</sup>/H<sup>+</sup> antiporter activity in hamster embryos is activated during fertilization. *Dev Biol.* 1999;208(1):244–252.
  13. Babcock DF, Rufo GA, Jr, Lardy HA. Potassium-dependent increases in cytosolic pH stimulate metabolism and motility of mammalian sperm. *Proc Natl Acad Sci USA.* 1983;80(5):1327–31.
  14. Babcock DF, Pfeiffer DR. Independent elevation of cytosolic [Ca<sup>2+</sup>] and pH of mammalian sperm by voltage-dependent and pH-sensitive mechanisms. *J Biol Chem.* 1987;262(31):15041–7.
  15. Marquez B, Suarez SS. Bovine sperm hyperactivation is promoted by alkaline-stimulated Ca<sup>2+</sup> influx. *Biol Reprod.* 2007;76(4):660–5.
  16. Leclerc C, Becker D, Buehr M, Warner A. Low intracellular pH is involved in the early embryonic death of DDK mouse eggs fertilized by alien sperm. *Dev Dyn.* 1994;200(3):257–67. [\[PubMed\]](#)
  17. Zhao Y, Chauvet PJ, Alper SL, Baltz JM. Expression and function of bicarbonate/chloride exchangers in the preimplantation mouse embryo. *J Biol Chem.* 1995;270(41):24428–34.
  18. Lane M, Baltz JM, Bavister BD. Na<sup>+</sup>/H<sup>+</sup> antiporter activity in hamster embryos is activated during fertilization. *Dev Biol.* 1999;208(1):244–52.
  19. Lane M, Bavister BD. Regulation of intracellular pH in bovine oocytes and cleavage stage embryos. *Mol Reprod Dev.* 1999;54(4):396–401.
  20. Lane M, Lyons EA, Bavister BD. Cryopreservation reduces the ability of hamster 2-cell embryos to regulate intracellular pH. *Hum Reprod.* 2000;15(2):389–94.
  21. Edwards LJ, Williams DA, Gardner DK. Intracellular pH of the preimplantation mouse embryo: effects of extracellular pH and weak acids. *Mol Reprod Dev.* 1998;50(4):434–42.
  22. Squirrell JM, Lane M, Bavister BD. Altering intracellular pH disrupts development and cellular organization in preimplantation hamster embryos. *Biol Reprod.* 2001;64(6):1845–54.
  23. Zander-Fox D, Mitchell M, Thompson JG, Lane M. Alterations in mouse embryo intracellular pH by DMO during culture impair implantation and fetal growth. *Reprod Biomed Online.* 2010;21(2):219–29.
  24. Fitzharris G, Baltz JM. Granulosa cells regulate intracellular pH of the murine growing oocyte via gap junctions: development of independent homeostasis during oocyte growth. *Development.* 2006;133(4):591–9.
  25. FitzHarris G, Siyanov V, Baltz JM. Granulosa cells regulate oocyte intracellular pH against acidosis in preantral follicles by multiple mechanisms. *Development.* 2007;134(23):4283–95.
  26. Phillips KP, Baltz JM. Intracellular pH regulation by HCO<sub>3</sub><sup>-</sup>/Cl<sup>-</sup> exchange is activated during early mouse zygote development. *Dev Biol.* 1999;208(2):392–405.
  27. Ferguson WJ, Braunschweiger KI, Braunschweiger WR, Smith JR, McCormick JJ, Wasemann CC, Jarvis NP, Bell DH, Good NE. Hydrogen ion buffers for biological research. *Anal Biochem.* 1980;104(2):300–10.
  28. Eagle H. Buffer combinations for mammalian cell culture. *Science.* 1971;174(8):500–3. [\[PubMed\]](#)
  29. Good NE, Winget GD, Winter W, Connolly TN, Izawa S, Singh RM. Hydrogen ion buffers for biological research. *Biochemistry.* 1966;5(2):467–77.