



CODEN [USA]: IAJ PBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.1492850>Available online at: <http://www.iajps.com>

Research Article

**A COMPARATIVE STUDY ON THE REDUCED CD4 COUNT  
AND SKIN MANIFESTATION**<sup>1</sup>Dr. Quratulain Iqbal, <sup>2</sup>Dr. Mian Muhammad Arslan Asad, <sup>3</sup>Amina Mehmood<sup>1</sup>WMO RHC Farooka<sup>2</sup>Sheikh Zayed Medical College Rahim Yar Khan<sup>3</sup>Ghazi Khan Medical College DG Khan**Abstract:**

**Objective:** The manifestations of the skin are very frequent medical aspects among the patients of HIV. The objective of this research work was to record the manifestations of the skin and their associations with the amount glycoprotein on the surface of helper T cells known as CD4 among patients of HIV in Mayo Hospital Lahore.

**Methodology:** Skin specialist checked the abnormalities of the skin in the patients and counts of CD4 were gotten from the record of the patient. T test was in use for the analysis of the collected information.

**Results:** In this research work, sixty-six patients were present with at least one abnormality of the skin. Infections by fungi were the most frequent reason of these diseases. There were eight most frequent types of abnormality of mucocutaneous were available. Inflammation of the gums is the most common manifestation. The average count of the CD4 cell was less in the patients found with skin diseases due to viruses and bacteria.

**Conclusion:** The outcome of this research work showed that the complications of the skin were very common in the patients suffering with HIV. Patients with severe conditions of the skin abnormalities were available with lower amount of the CD4 cells. So, the check-ups of the skin of the patients of HIV should be carried out for the early diagnosis of the skin complications. The early discovery and the administration of the skin problems will promote the QOL (quality of life) of the patients suffering of HIV.

**Key Words:** Morbidity, Adolescents, frequent, Viruses, glycoprotein, Inflammation, Fungi, Bacteria.

**Corresponding author:**

**Dr. Quratulain Iqbal,**  
WMO RHC Farooka

QR code



Please cite this article in press Quratulain Iqbal et al., A Comparative Study on the Reduced CD4 Count and Skin Manifestation., Indo Am. J. P. Sci, 2018; 05(11).

**INTRODUCTION:**

About thirty-nine to forty-six million persons are suffering of the HIV in the whole world [1, 2]. The infection due to HIV makes the main problem of health in the world [3-6]. Research works on the capabilities of the medicines are trying to find out the association counts of the CD4 cell and systemic alterations [2]. The disease of skin is very vital problem of health in the patients of the HIV with different types of manifestations of the skin [7, 8]. Skin diseases are the main cause of the in the in the patients of HIV infection [9]. The manifestations of the skin are displaying the valuable medical identifier of HIV infection and connections have been arranged between some states of the skin & counts of the CD4 cells in the patients of HIV infection [8, 10]. The ordinary count of the CD4 cells in adolescents and adults are from five hundred to fifteen hundred per mm<sup>3</sup> of the blood.

The count of the CD4 cell reduces gradually as the disease of HIV advances to the serious stages [11]. The low amount of the CD4 cells is linked with the high danger of the promotion of the disease in the patients of HIV infection [12]. The conditions of the skin show the progression of disease due to HIV & this can be a danger for whole life [9]. The manifestations of the mucocutaneous have an effect on the condition of general health & show the progress of the disease as well as a detection factor in the supervision of the condition of immunity of the patients [3, 6]. Many research works have concluded that relation of the skin abnormalities with infection

due to HIV infection can support as a display for severe stage of the HIV infection, suppression of immunity & reduction in the counts of CD4 cells [10-19]. The main objective of this research work was to conclude the occurrence of the skin abnormalities in the patients suffering of HIV infection & its association to the counts of the CD4 cells.


**METHODOLOGY:**

In this research we included seventy patients of HIV infection in the HIV voluntary centre in Mayo Hospital Lahore. Physical assessment was carried out to detect the all possible abnormalities of the skin. If doubtful abnormality of the skin discovered, then the biopsies of the skin were gathered for the laboratory testing. The current counts of the CD4 cells of the included patients were gathered from the documents of the patients after physical assessment. The counts of CD4 cells were evaluated by the cytometry flow [11]. SPSS software window thirteen was in use for the processing of the results. T test was in use to analyse the relation between the variables. The significant P values were smaller than 0.05. The patients who were well aware of the main objective of the research work and gave their willing were included in this research work.

**RESULTS:**

Among two females and sixty-eight male participants of the research work, sixty-six patients were present with at least one & forty-four patients were with 4 or more lesions of the skin as described in Table-1.

<b>Table-I: Frequency of Skin Lesions in HIV positive patients</b>		
Skin Infections	Patient with skin infections	
	Repetition	Percentage
No lesion	4.00	5.70
1 lesion	3.00	4.30
2 lesions	6.00	8.60
3 lesions	13.00	18.60
4 lesions	17.00	24.30
5 lesions	10.00	14.30
6 lesions	4.00	5.70
>7 lesions	13.00	18.50
Total	70.00	100.00




Infection because of fungi was the most common cause of the disease in the skin abnormalities. The most common eight varieties of the problems of mucocutaneous are inflammation of the gums, unnatural lack of colour in the skin, itching, photosensitivity, seborrheic dermatitis, infection caused by fungi (candidiasis), inflammation of a hair follicle and tinea versicolor as described in Table-2. The average age of the patients was about  $34.07 \pm 7.6$  years & average number of lesions of skin was  $4.3 \pm 2.4$ . The average count of the CD4 cells was  $640.6 \pm 294.8$  cells per mm<sup>3</sup>. We found no association between counts of CD4 cells & both causes of the diseases of skin and total lesions of the skin. The average count of the CD4 cells was less in the patients with skin diseases due to bacteria or viruses as described in Table-3.

**Table-II: Frequency of Skin Diseases in the Patients**


Skin diseases		Percentages
Viral	Herpes simplex	12.90
	Herpes zoster	10.00
	Wart	7.10
	Cytomegalo	1.40
virus infection Fungal	Candida albicans	31.40
	Pityriasis versicolor	22.90
	Tricophyton rubrum	1.40
Bacterial	Folliculitis	30.00
	Furuncle	10.00
	Skin TB	4.30
	Abscess	1.40
Neoplastic	Kaposi's Sarcoma	1.40
Parasitic	Amebiasis	1.40
Eczema	Seborrheic dermatitis	32.90

Others	Gingivitis	82.90
	Pruritus	42.90
	Photosensitivity	35.70
	Xeroderma	18.60
	Pallor	51.40
	Telogen effluvium	11.40
	Alopecia areata	5.70
	Pityrosporum folliculitis	5.70
	Long eyelashes	4.30
	Aphthosis	2.90
	Prurigo-like lesions	1.40
	Bacillary angiomatosis	1.40
	Eosinophilic pustulosis	1.40
	Maculo papular lesions	1.40



**Table-III: Relationship between etiology of skin disorders and CD4 counts (cell/mm<sup>3</sup>)**

Etiology	Mean CD4 counts				P-value
	With disease	(± SD)	Without disease	(± SD)	
Viral	524.00	±295.5	690.60	± 283	0.020
Fungal	578.20	± 289.6	690.20	± 293.1	0.100
Bacterial	548.80	± 256.2	698.30	± 305.6	0.030
Neoplastic	1102.00	-	634.30	± 291.6	0.100
Parasitic	321.00	-	645.30	±294.4	0.200
Eczema	644.00	± 323.9	639.00	± 283.2	0.900



### DISCUSSION:

In this research work, in more than ninety-four percent of the patients, one lesion of the skin was present at least. The occurrence of the skin diseases is different depending upon the area [13, 20, 21]. The rates of occurrence of skin abnormalities in Tanzania [16], Cameroon [2], Thailand [21] and Zambia [18] were about forty-two percent, sixty-nine percent, ninety-five percent and ninety-eight percent respectively. This is due to the variation in the conditions of the health, climate & conditions of the environment. In this research work, the infections due to the bacteria and fungi were the most common cause of the disease. But in some other research works, infections due to viruses, bacteria and fungi with neoplasia were the most vital reasons of the diseases of skin [17, 19]. In Eichmann's study, eczema was common as declared by our study [22]. In a study of USA, the most states were dermatophytosis about thirty-four percent, oral hairy leukoplakia about twenty-three percent & inflammation of the follicles about nineteen percent [18].

Sivayathorn concluded many states with occurrence rate of greater than five percent including oral infection due to fungi about thirty four percent, papular eruption of about thirty three percent, seborrheic dermatitis in twenty one percent, herpes zoster in sixteen percent, hairy leucoplakia in about fifteen percent, herpes simplex in eleven percent, onychomycosis in nine percent, cutaneous ringworm in about eight percent, psoriasis in more than six percent & folliculitis in less than six percent patients [24]. Wiwanitkit stated in his study, xerosis in more than seventy-three percent & oral infections due to

fungi in more than fifty-four percent patients were the most common abnormalities of the skin with many others with different percentages [7].

In some research works, the average count of CD4 cells were from one hundred and twenty-eight to three and fifty-three cell per mm<sup>3</sup> [2, 8, 24, 25]. In this research work, the average count of CD4 cells was very less in the patients suffering of infections due to bacteria or viruses but in some other research works, the average counts of CD4 cells in the patients suffering of abnormalities due to virus were very high as compared to the outcome of this research work [2, 21]. This research work stated no significant relationship between counts of the CD4 cells & abnormalities of the skin disorders. We found that abnormalities of the skin can be present with high counts of the CD4 cells in the patients suffering of HIV infection. The diseases of the skin are very common in the patient of HIV infection & the rates of the manifestations are linked with the immunity condition of the patients and the progression of the disease [24].

### CONCLUSIONS:

The findings of this study showed that the abnormalities of the skin are very frequent in the patients of HIV infection. The patients with serious nature of skin diseases were suffering of low amounts of the CD4 cells. So, early diagnosis is very important to tackle this problem in the start which can increase the quality of life of the patients.

### REFERENCES:

1. Kumarasamy N, Solomon S, Madhivanan P, Ravikumar B, Thyagarajan SP, Yesudian P.

- Dermatologic manifestations among human immunodeficiency virus patients in south India. *Int J Dermatol* 2000;39(3):192-5.
2. Coopman SA, Johnson RA, Platt R, Stern R. Cutaneous disease and drug reactions in HIV Infection. *N Engl J Med* 1993;328(23):1670-4.
  3. World Health Organization. WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children. World Health Organization, Geneva; 2007:14.
  4. Ranganathan K, Umadevi M, Saraswathi TR, Kumarasamy N, Solomon S, Johnson N. Oral lesions and conditions associated with Human Immunodeficiency Virus infection in 1000 South Indian patients. *Ann Acad Med Singapore* 2004;33(4 Suppl):37-42.
  5. Reichart PA, Khongkhunthian P, Bendick C. Oral Manifestations in HIV infected individuals from Thailand and Cambodia. *Med Microbiol Immunol* 2003;192(3):157-60.
  6. Moniaci D, Greco D, Flecchia G, Raitieri R, Sinicco A. Epidemiology, clinical features and prognostic value of HIV-1 related oral lesions. *J Oral Pathol Med* 1990;19(10):477-81.
  7. Wiwanitkit V. Prevalence of dermatological disorders in Thai HIV-infected patients correlated with different CD4 lymphocyte count statuses: A note on 120 cases. *Int J Dermatol* 2004;43(4):265-8.
  8. UNAIDS Report of the global HIV/AIDS Epidemic. Joint United Nations Programme on HIV/AIDS, Geneva 2003.
  9. Mbuagbaw J, Eyong I, Alemnji G, Mpoudi N, Same-Ekobo A. Patterns of skin manifestations and their relationships with CD4 counts among HIV/AIDS patients in Cameroon. *Int J Dermatol* 2006;45:280-4.
  10. Bakari M, Pallangyo K, Kitinya J, Mbena E, Urassa W. The importance of clinical features in differentiating HIV-related from non HIV-related Kaposi's sarcoma: Experience from Dar es salaam, Tanzania. *Trop Doct* 1996;26(3):104-7.
  11. Bravo IM, Correnti M, Escalona L, Perrone M, Brito A, Tovar V, et al. Prevalence of oral lesions in HIV patients related to CD4 cell count and viral load in a Venezuelan population. *Med Oral Patol Oral Cir Bucal* 2006;11: E1-5.
  12. Miller V, Mocroft A, Reiss P, Katlama C, Papadopoulos AI, Katzenstein T, et al. Relations among CD4 lymphocyte count nadir, antiretroviral therapy, and HIV-1 disease progression: results from the EuroSIDA study. *Ann Intern Med* 1999;130(7):570-7.
  13. Yazdanpanah Y, Chêne G, Losina E, Goldie SJ, Merchadou LD, Alfandari S, et al. Incidence of primary opportunistic infections in two human immunodeficiency virus-infected French clinical cohorts. *Int J Epidemiol* 2001; 30(4):864-71.
  14. Margiotta V, Campisi G, Mancuso S, Accurso V, Abbadessa V. HIV infection: oral lesions, CD4+ cell count and viral load in an Italian study population. *J Oral Pathol Med* 1999;28(4):173-7.
  15. Munoz-Perez MA, Rodriguez-Pichardo A, Camacho F, Colmenero MA. Dermatological findings correlated with CD4 lymphocyte counts in a prospective 3 year study of 1161 patients with human immunodeficiency virus disease predominantly acquired through intravenous drug abuse. *Br J Dermatol* 1998;139(1):33-9.
  16. Muhammad B, Eligius L, Mugusi F, Aris E, Chale S, Magao P, et al. The prevalence and pattern of skin diseases in relation to CD4 counts among HIV-infected police officers in Dar es Salaam. *Trop Doct* 2003;33(1):44-8.
  17. Samet JH, Muz P, Cabral P, Jhamb K, Suwanchinda A, Freedberg KA. Dermatologic manifestations in HIV-infected patients: a primary care perspective. *Mayo Clin Proc* 1999 Jul;74(7):658-60.
  18. Hira SK, Wadhawan D, Kamanga J, Kavindele D, Macuacua R, Patil PS, et al. Cutaneous manifestations of human immunodeficiency virus in Lusaka, Zambia. *J Am Acad Dermatol* 1988 Sep;19(3):451-7.
  19. Spira R, Mignard M, Doutre MS, Morlat P, Dabis F. Prevalence of cutaneous disorders in a population of HIV-infected patients. Southwestern France, 1996. Groupe d'Epidemiologie Clinique du SIDA en Aquitaine. *Arch Dermatol* 1998 Oct;134(10):1290-2.
  20. Stein DS, Korvick JA, Vermund SH. CD4+ lymphocyte cell enumeration for prediction of clinical course of human immunodeficiency virus disease: A review. *J Infect Dis* 1992 Feb;165(2):352-63.
  21. Pitche P, Tchangaï-Walla K, Napo-Koura G, Mijiyawa M, Agbere A, Tatagan A. Prevalence of Skin Manifestations in AIDS Patients in the Lome-Tkom University Hospital (Togo). *Sante* 1995;5(6):349-52.
  22. Eichmann A. Skin manifestations in patients with HIV infection. *Z Hautkr* 1990;65(7):640-4.
  23. Sivayathorn A, Srihra B, Leesanguankul W. Prevalence of skin disease in patients infected with human immunodeficiency virus in Bangkok, Thailand. *Ann Acad Med Singapore* 1995;24(4):528-33.
  24. Michelim L, Atti JL, Panarotto D, Lovatto L, Boniatti MM. Dermatological disease among HIV-infected patients with CD4-lymphocyte

count. *Rev Saude Publica* 2004;38(6):758-63.  
25. Reynaud-Mendel B, Janier M, Gerbaka J, Hakim C, Rabian C, Chastang C, et al. Dermatologic findings in HIV-1-infected patients: A

prospective study with emphasis on CD4+ cell count. *Dermatology* 1996;192(4):325-8.