



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**Available online at: <http://www.iajps.com>

Research Article

**FORMULATION AND EVALUATION OF BARICITINIB
TRANSFERSOMES****Prashanti Chitrapu¹, Sukanya Mandala², D. Suchitra³, Konda Shravan Kumar⁴**^{1,2}Dept. of Pharmaceutical Chemistry, Gyana Jyothi College of Pharmacy, Telangana³Dept. of Pharmacy, University college of technology, Osmania University, Telangana⁴Dept. of Pharmaceutics, Trinity College of Pharmaceutical Sciences, Peddapalli**Abstract:**

Transfersomes are specially optimized particles or vesicles, which can respond to an external stress by rapid and energetically inexpensive, shape transformations. The present study was carried out to use statistical techniques of quality by design to develop and optimize Methotrxate Transfersomes by conventional rotary evaporation sonication techniques. A 3² full factorial design was used to understand the main effects and interaction of formulation variables. The relationship between the dependent and independent variables was further elucidated using contour plots. Then, experimental design combined with desirability functions to predict the desired quality.

A conventional rotary evaporation sonication technique was developed to prepare Baricitinib transfersomes using Phospholipon 90 G as a lipid (SPC) and SPAN 80 as an edge activator (EA). Preformulation studies proved the purity of drug and compatibility of drug and excipients was evaluated by DSC study. BCT loaded transfersomes was successfully formulated using 3² full factorial design and desirability function. The % entrapment efficiency, deformability index and particle size were highly dependent on the SPC: EA ratio, and BCT concentration for the preparation of BCT loaded Transfersomes. SPC:EA ratio and BCT concentration had a positive effect on % Entrapment efficiency and negative effect on deformability. But particle size was not much affected. Using contour plots, response surface plots and % Bias, formulation was optimized. Batch M5 (1:1 PC: EA, 2% w/v BCT) is best batch among nine combinations. Surface morphology of transfersomes was evaluated by TEM study. A good interaction between drug and excipients were found with FTIR study of transfersomes. It was advisable to store BCT-TFS at refrigerated conditions (4±2 °C). Optimum drug: lipid ratio is 0.1-0.4. BCT-TFS was successfully optimized with experimental design and desirability function.

Keywords: Baricitinib, transfersomes, BCT-TFS.

Corresponding author:**Prashanti Chitrapu,**

Dept. of Pharmaceutical Chemistry,

Gyana Jyothi College of Pharmacy, Telangana

prashantichitrapu@yahoo.com

QR code



Please cite this article in press Prashanti Chitrapu et al *Formulation And Evaluation Of Baricitinib Transfersomes*, Indo Am. J. P. Sci, 2018; 05(04).

INTRODUCTION:

Transfersome (TFS) is a term registered as a trademark by the German company IDEAAAG, and used by it to refer to its proprietary drug delivery technology. The name means “carrying body”, and is derived from the Latin word ‘transferre’, meaning to carry across”, and the Greek word „soma“, for a „body“. A transfersome carrier is an artificial vesicle designed to be like a cell vesicle or a cell engaged in exocytosis, and thus suitable for controlled and, potentially targeted, drug delivery. Transfersome is a highly adaptable and stress-responsive, complex aggregate. Its preferred form is an ultra-deformable vesicle possessing an aqueous core surrounded by the complex lipid bilayer. Interdependency of local composition and shape of the bilayer makes the vesicle both self-regulating and self-optimizing. This enables the transfersome to cross various transport barriers efficiently, and then act as a drug carrier for non-invasive targeted drug delivery and sustained release of therapeutic agents. The presence of surface-active agents in the transfersomes enhances the rheological properties and sensitivity to the driving force which results from water concentration gradient across the skin. This enhances the propensity of sufficiently large but deformable pentrant, transfersomes to move across the skin barrier. Such capability combined with the inclination to deform into elongated shapes while maintaining the vehicle integrity can explain the usually high efficiency of transfersomes across the skin.

Preparation of transfersomes has many variables which significantly affect the characteristics of it. Among all variable 3 variables were significant as described by results of placket burman design. The present study, therefore, deals with the optimization of formulation variables to design the best product under conditions of

competitive objectives, because interactive effects via a trial-and-error approach are time-consuming and often unsuccessful. Mathematical optimization by means of an experimental design is most helpful in shortening the experimental time⁶.

MATERIALS AND METHODS:

Materials

Baricitinib was received as gift sample from West Coast Pharmaceutical Pvt. Ltd. Ahmedabad, (India). Phospholipon 90 G (lipid) was obtained as a gift sample from Lipoid AG, Germany. Double distilled water was prepared in laboratory for study. All materials used for study conformed to USP 24 standards and purchased from ACS chemical Pvt. Ltd. Ahmedabad (India).

METHODS

Preparation of Baricitinib Transfersomes

Baricitinib Transfersomes were prepared by conventional rotary evaporation sonication method. Precisely, Phospholipon 90 G (SPC) mixed with Edge activator Span 80 were taken in a clean, dry, round bottom flask and the lipid mixture (100 mg/ml) was dissolved in Methanol: Chloroform (1:2). The organic solvent was removed by rotary evaporator above the lipid transition (40°C). Final traces of solvent were removed under vacuum overnight. The deposited lipid film was hydrated with 15 ml PBS (pH 7.4) containing Drug Baricitinib to furnish the desired concentration in the final preparation by rotation at 50 rpm for 1 h at 50 °C. The resulting vesicles were swollen for 2 hr at room temperature to get large multilamellar vesicles (LMLVs). The thick suspension thus obtained was broken by sonication for 30 min at 4 °C at a frequency of 53 kHz to achieve desired vesicle size (200-300 nm)⁹. Composition of Baricitinib transfersomes were described in Table 1.

Table 1: Composition of 3² factorial design of BCT-TFS

Coded factors	Factors/Levels	-1	0	1
X1	SPC: EA	0.5:1	1:1	2:1
X2	BCT	1% w/v	2% w/v	3% w/v

Optimization of formulation parameters

Optimization of formulation was carried out using Experimental design and desirability function. A two-factor, three-level full factorial design was applied for the optimization procedure using DOE++ software (Version 1.0.7.2 ID –DS-1, ReliaSoft Corporation, USA). The ratio of SPC: EA and Baricitinib (BCT) concentration were used to prepare each of the 9 formulations are given in Table 6.2. These high,

medium, and low levels were selected from the preliminary experimentation. After generating the polynomial equations, relating the dependent and independent variables, the process was optimized for the % Entrapment efficiency (Y1), Deformability Index (Y2), Particle size (Y3). After the fitting of the mathematical model, the desirability function was used for the optimization.

Table 2: Formulation of 3² factorial design batches of BCT-TFS

Run	Coded value		Actual value				
	X1	X2	SPC: EA	SPC in gm	EA in gm	BCT in %w/v	BCT in gm
M1	-1	-1	0.5:1	0.5	1.0	1	0.15
M2	-1	0	0.5:1	0.5	1.0	2	0.30
M3	-1	1	0.5:1	0.5	1.0	3	0.45
M4	0	-1	1:1	0.75	0.75	1	0.15
M5	0	0	1:1	0.75	0.75	2	0.30
M6	0	1	1:1	0.75	0.75	3	0.45
M7	1	-1	2:1	1.0	0.5	1	0.15
M8	1	0	2:1	1.0	0.5	2	0.30
M9	1	1	2:1	1.0	0.5	3	0.45

Characterization of Baricitinib transfersomes Determination of percent drug content

%Entrapment Efficiency

Transfersome entrapped Baricitinib was estimated by centrifugation method. The prepared transfersome were placed in centrifugation tube and centrifuged at 14000 rpm for 30 minute. The supernatant (1ml) was withdrawn and diluted with phosphate buffer (pH 7.4). The untrapped Baricitinib was determined by UV spectrophotometer at 305.2 nm.

%Entrapment Efficiency = (Total Drug-Untrapped drug)/Total Drug X 100

Deformability Index (Vesicle Elasticity Measurement)

The deformability study was done for the transfersomal formulation using a home- built device¹⁴. The elasticity of transfersomes vesicles were measured by extrusion method. The transfersomes formulation were extruded through filter membrane (pore size diameter 100 nm), using a stainless steel filter holder (50 mm diameter), by applying a pressure of 2.5 bar. The quantity of vesicles suspension extruded in 5 minutes was measured. The deformability index of prepared batches is depicted in Table 6.4.

$$E = J * (rv / rp)^2$$

Where, E = Elasticity of vesicles membrane,
J = Amount of suspension extruded in 5 minutes,
rv = Vesicles size,
rp = pore diameter.

Particle Size and Surface Charge

The droplet size and zeta potential of the transfersomes was determined by a Malvern Particle sizer and Zeta Potential Analyzer (Malvern Instruments Ltd., UK) at room temperature¹⁵. 1 ml of the transfersome suspension was diluted with deionized water.

Transmission Electron Microscopy

Transmission Electron Microscopy (TEM) was used to visualize the transfersomal vesicles using TEM microscope (Model: Philips Tecnai 20 G2, Holland) at SICART, Vallabh Vidhynagar. The vesicles were dried on a copper grid and adsorbed with filterpaper¹⁵. After drying, the sample was viewed under the microscope at 10–100 k magnification at an accelerating voltage of 100 kV. The TEM image of best batch is shown in Figure.

Fourier Transform Infrared Spectroscopy

Fourier transform infrared spectroscopy (FTIR) study was carried out to confirm structure of formulation. FTIR spectra of pure drug, lipid and formulated Transfersomes containing drug were recorded on FTIR Spectrophotometer (SHIMADZU, Japan) at Ratnamani Health care, Indrad. The scanning range

was from 4000 to 600 cm^{-1} and the resolution was 1 cm^{-1} .

Physical stability

This study was carried out in stability chambers at Ratnamni health care. After measuring the initial percentage entrapment of the drug in the various formulations, the three batches of the same formulation were stored in sealed glass ampoules at different temperature⁹. According to ICH, physical stability was carried out at refrigeration temperature

(4 ± 2 °C) and room temperature (25 ± 2 °C/65% RH \pm 5% RH) for a period of 3 months

RESULTS AND DISCUSSION:

Melting point determination

DSC thermogram of Baricitinib explains that melting point of drug was 192.87 °C (Figure 1). Reported range of melting point of BCT is 185-204°C¹⁷⁻¹⁸. So BCT has confirmed its physicochemical property.




Figure 1: DSC thermogram of Baricitinib

Drug-Excipients compatibility study

Compatibility of drug and lipid was confirmed from comparison of DSC thermogram of Baricitinib and physical mixture of Baricitinib and lipid (Figure 2).




Figure 2: DSC spectra of mixture of Baricitinib and lipid

%Entrapment efficiency

The maximum entrapment efficiency 90.71 ± 3.4 was found in case of transfersomal formulation M4 (Table 3). Lipid composition of the batch M4 was higher in comparison to that of other batches and this may result in higher entrapment efficiency measured with transfersomes. Further, transfersomes contains a mixture of lipid and membrane softener, SPAN 80. The lipid is a stabilizing factor and SPAN 80 is a destabilizing factor. In the preparation method of transfersomes, the vesicles content is exchanged with the dispersion medium during breaking and resealing of phospholipid bilayer as they are sonicated using probe sonicator. During the successive cycles, the drug stayed inside the transfersomes suggesting that interaction between lipid membrane and drug did not allow free displacement of drug.

Table 3: Results of dependent variables of 3² factorial design batches of BCT-TFS

Run	Dependent variables		
	Y1	Y2	Y3
M1	89.02±1.48	38.21±1.23	265.8±1.2
M2	64.67±0.46	32.87±1.14	272.5±1.8
M3	45.92±0.48	33.50±2.09	325.4±0.7
M4	90.71±0.66	21.77±0.70	267.3±2.4
M5	84.77±0.26	27.13±1.13	314.2±1.4
M6	58.51±0.35	26.41±1.06	317.5±1.5
M7	90.45±0.66	17.63±1.99	281.8±0.5
M8	88.32±0.40	10.65±0.64	332.5±2.2
M9	73.21±0.3	14.82±1.7	342.4±1.

Particle size and surface charge

Droplet size was measured by Malvern particle analyzer. Average sizes of all transfersomes formulations were in the range of 250-350 nm . This suggests that prepared transfersomes are good for penetration through skin as they have nano sizes .

Differences in particle size within and among the batches were observed to be insignificant ($P < 0.05$). It was also noticed that size of BCT-entrapped transfersomes increased with the increasing SPC: EA ratio. Increase in the concentration of surfactant was found to decrease in BCT-TFS size significantly ($P < 0.05$).




Figure 3: Particle size distribution of transfersomes batch M5

Transmission electron microscopy

TEM image for optimized Transfersomes is shown with SPC: EA ratio; 1:1 and BCT concentration: 2% w/v (Figure 6.14). Transmission electron microscopy was used to characterize transfersomes. These carriers invariably appeared as unilamellar vesicles. Bioactive molecules larger than 500 Da normally do not cross the skin. This prevents non-invasive delivery of high-molecular-weight bioactive. In order to cross the intact mammalian skin, transfersomes should be capable of passing through pores of diameter less than 50 nm under influence of suitable transdermal gradient: FTIR study




Figure 4: Transmission electron micrograph of optimized BCT-TFS batch M5

The Fourier transform infrared spectroscopy (FTIR) spectrum of the formulation (transfersomes) was compared with pure drug and lipid spectra. This study reveals that all major lipid peaks were observed in spectra of Baricitinib transfersomes and Baricitinib peaks were merged and shifted in the same. So it was concluded that drug was completely incorporated in transfersomes.




Figure 5: IR spectra of Baricitinib transfersomes

Physical stability of Baricitinib Transfersomes

Stability is an essential quality attribute for drug products. This evaluation checked by pharmaceutical scientist and regulators quantify drug product stability and shelf life. Drug stability concerns about drug product safety, efficacy, and quality, found it to appropriate³⁰. Physical stability represents the ability of a product to maintain its physical dimensions and properties when exposed to conditions normally encountered in its service environment³¹. Figure 6.18 explains that % drug loss from transfersomes was higher at room temperature (25 ± 2 °C/ $65\% \text{ RH} \pm 5\% \text{ RH}$) as compared to refrigeration temperature (4 ± 2 °C).




Figure 6: Physical stability of BCT-TFS Batch M5

Derivation of Optimum Drug: Lipid ratio

From the data of 3^2 factorial design, drug:lipid ratio can also take for desired drug loading. Drug:lipid ratio is derived and drug loading (%EE) is compared. Table 6.8 explains as drug:lipid ratio is increased from 0.1 to 0.9, drug loading capacity of transfersomes is decreased from 90% to 40%. It is recommended from the results that optimum drug:lipid ratio can be used is 0.1-0.4.

Table 6.8: Optimization of Drug:Lipid ratio

Run	SPC	EA	BCT	BCT:SPC	%Drug loading
M7	1	0.5	0.15	0.15	90.45±0.66
M4	0.75	0.75	0.15	0.2	90.71±0.66
M1	0.5	1	0.15	0.3	89.02±1.48
M8	1	0.5	0.3	0.3	88.32±0.40
M5	0.75	0.75	0.3	0.4	84.77±0.26
M9	1	0.5	0.45	0.45	73.21±0.35
M2	0.5	1	0.3	0.6	64.67±0.46
M6	0.75	0.75	0.45	0.6	58.51±0.35
M3	0.5	1	0.45	0.9	45.92±0.48

CONCLUSION:

Transfersomes are specially optimized particles or vesicles, which can respond to an external stress by rapid and energetically inexpensive, shape transformations. A conventional rotary evaporation sonication technique was developed to prepare Baricitinib transfersomes using Phospholipon 90 G as a lipid (SPC) and SPAN 80 as an edge activator (EA). Preformulation studies proved the purity of drug and compatibility of drug and excipients was evaluated by DSC study. BCT loaded Transfersomes was successfully formulated using 3² full factorial design and desirability function. The % entrapment efficiency, deformability index and particle size were highly dependent on the SPC: EA ratio, and BCT concentration for the preparation of BCT loaded Transfersomes. SPC:EA ratio and BCT concentration had a positive effect on % entrapment efficiency and negative effect on deformability. But particle size was not much affected. Using contour plots, response surface plots and % Bias, formulation was optimized. Batch M5 (1:1 PC: EA, 2% w/v BCT) is best batch among nine combinations. Surface morphology of transfersomes was evaluated by TEM study. A good interaction between drug and excipients were found with FTIR study of transfersomes. It was advisable to store BCT-TFS at refrigerated conditions (4±2 °C). Optimum drug:lipid ratio is 0.1-0.4. BCT-TFS was successfully optimized with experimental design and desirability function.

REFERENCES

1. Carol E. Rheumatoid Arthritis Treatments - 2008 ACR Recommendations [Internet]. About.com; 2008 [cited 2013 Jul 25]. Available from: http://arthritis.about.com/od/rheumatoidarthritis/a/RA_treatments.htm.
2. Colebatch AN, Marks JL, Edwards CJ. Safety on non-steroidal anti-inflammatory drugs, including aspirin & paracetamol in people receiving Baricitinib for inflammatory arthritis. *Cochrane Database of Systemic Reviews*. 2011; 9(1): CD008872. DOI: 10.1002/14651858.
3. Walve JR, Bakliwal SR, Rane BR, Pawar SP. Transfersomes: a surrogated carrier for transdermal drug Delivery system. *International journal of applied biology and pharmaceutical technology*. 2011; 2(1): 204-213.
4. Dua K, Pabreja K, Sharma VK, Singh Sara UV, Agrawal DK. Transfersomes: pivotal role in drug delivery. *Ethiop. J. Health Biomed. Sci*. 2009; 2(1): 47-52.
5. Prajapati ST, Patel CG, Patel CN. Transfersomes: A vesicular carrier system for transdermal drug

delivery. *Asian Journal of Biochemical and Pharmaceutical Research*. 2011; 1(2): 507-524.

6. Patel RB, Parikh RH. Preparation and formulation of transfersomes containing an antifungal agent for transdermal delivery: Application of Plackett-Burman design to identify significant factors influencing vesicle size. *J. Pharm. Bioall. Sci*. 2012; 4: 60-61. DOI:10.4103/0975-7406.94140.
7. Garg R, Gupta GD, Kaur B. Preformulation: a need for dosage form design. *Pharmaceutical formulation development*. 2008; 6(1). Available from: <http://www.pharmainfo.net/reviews/preformulation-on-need-dosage-form-design>.
8. Stability testing of new drug substances and products, Q1A(R2), ICH guideline for stability testing, 6 February, 2003; 3-4.
9. Maurya SD, Agrawal S, Maurya G. Enhanced transdermal delivery of indinavir sulfate via Transfersomes. *International journal of comprehensive pharmacy*. 2010;1(6): 1-7.
10. Shahinaze A, Fouad, Emad B, Basalious, Mohamed A, El-Nabarawi, Saadia A, Tayel. Microemulsion and poloxamer microemulsion-based gel for sustained transdermal delivery of diclofenac epolamine using in-skin drug depot: In vitro/in vivo evaluation. *Int J Pharm*. 2013; 453(2):569-78.
11. Li F1, Song S, Guo Y, Zhao Q, Zhang X, Pan W, Yang X. Preparation and pharmacokinetics evaluation of oral self-emulsifying system for poorly water-soluble drug Lornoxicam. *Drug Deliv*. 2015; 22(4):487-98
12. Block, L.H., Pharmaceutical emulsions and microemulsions. *Pharmaceutical dosage forms, disperse systems*. 2001; 2: 47-109.
13. Tadros, T. Formation and stability of nano-emulsions. *Advances in colloid and interface science*. 2004;108:303-318.
14. Hongwu Sun, Kaiyun Liu, Wei Liu, Wenxiu Wang, Chunliang Guo, Bin Tang, Jiang Gu, Jinyong Zhang, Haibo Li, Xuhu Mao, Quanming Zou, and Hao Zeng. Development and characterization of a novel nanoemulsion drug-delivery system for potential application in oral delivery of protein drugs *Int J Nanomedicine*. 2012; 7: 5529-5543
15. Zheng Wang, Hong-Jie Mu, Xue-Mei Zhang, Peng-Kai Ma, Sheng-Nan Lian, Feng-Pu Zhang, Sheng-Ying Chu, Wen-Wen Zhang, Ai-Ping Wang, Wen-Yan Wang, Kao-Xiang. Lower irritation microemulsion-based rotigotine gel: formulation optimization and in vitro and in vivo studies. *Int J Nanomedicine*. 2015; 10: 633-644.
16. Azar Kajbafvala, Alireza Salabat, Anayatollah

- Salimi. Formulation, characterization and *in-vitro/ex-vivo* evaluation of quercetin-loaded microemulsion for topical application. *Pharmaceutical Development and Technology* 2016; 1-10
17. Maulvi FA, Desai AR, Choksi HH, Patil RJ, Ranch KM, Vyas BA, Shah DO. Effect of surfactant chain length on drug release kinetics from microemulsion-laden contact lenses. *Int J Pharm.* 2017; 524(1-2):193-204.
 18. Maulvi F, Desai A, Choksi H, Patil R, Ranch K, Vyas B, Shah D. Effect of surfactant chain length on drug release kinetics from microemulsion-laden contact lenses. *Int J Pharm.* 2017; 524(1-2):193-204.
 19. M.L De Martin, E.L Cussler, Predicting subjective spreadability, viscosity, and stickiness. *J. Pharm. Sci.*; 64: 976-982
 20. Shinde, Pokharkar S, Modani S. Design and evaluation of microemulsion gel system of nadifloxacin. *Indian J Pharm Sci.* 2012; 74(3):237-47
 21. Mrunali R. Patel, Rashmin B. Patel, Jolly R. Parikh, Bharat G. Patel. Novel isotretinoin microemulsion-based gel for targeted topical therapy of acne: formulation consideration, skin retention and skin irritation studies. *Appl Nanosci.* 2016; 6:539-553
 22. Abd-Allah FI, Dawaba HM, Ahmed AM. Preparation, characterization, and stability studies of piroxicam-loaded microemulsions in topical formulations *Drug Discov Ther.* 2010; 4(4):267-75.
 23. Mrunali R. Patel, Rashmin B. Patel, Jolly R. Parikh, Bharat G. Patel. Novel isotretinoin microemulsion-based gel for targeted topical therapy of acne: formulation consideration, skin retention and skin irritation studies. *Appl Nanosci.* 2016; 6:539-553.
 24. Patel HK, Barot BS, Parejiya PB. Topical delivery of clobetasol propionate loaded microemulsion based gel for effective treatment of vitiligo: ex vivo permeation and skin irritation studies. *Colloids Surf B* 2013; 102:86-94.
 25. Ngawhirunpat T, Worachun N, Opanasopit P. Cremophor RH40-PEG 400 microemulsions as transdermal drug delivery carrier for ketoprofen. *Pharm Dev Technol* 2013; 18:798-803.
 26. Tsai YH, Lee KF, Huang YB. In vitro permeation and in vivo whitening effect of topical hesperetin microemulsion delivery system. *Int J Pharm* 2010; 257-262.
 27. Ramesh Gannu, Vamshi Vishnu Yamsani, Shraavan Kumar Yamsani. Enhanced bioavailability of lacidipine via microemulsion based transdermal gels Formulation optimization, ex vivo and in vivo characterization. *International Journal of Pharmaceutics* 2010; 388 :231-241
 28. Ujwala Shinde,* Sharda Pokharkar, and Sheela Modani Design and Evaluation of Microemulsion Gel System of Nadifloxacin *Indian J Pharm Sci.* 2012; 74(3): 237-247
 29. Eskandar Moghimipour, Anayatollah Salimi, Masoud Karami, Sara Isazadeh Jundishapur. Preparation and Characterization of Dexamethasone Microemulsion Based on Pseudoternary Phase Diagram. *J Nat Pharm Prod.* 2013 Aug; 8(3): 105-112.
 30. Draize J, Woodard G, Calvery H. Methods for the study of irritation and toxicity of substances topically applied to skin and mucous membranes. *J Pharmacol Exp Ther.* 1944; 82: 377-390.