



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.1488011>Available online at: <http://www.iajps.com>

Research Article

**A STUDY ON RELATIONSHIP OF BLOOD PRESSURE AND  
HYPERTENSION IN THE RISK ASSESSMENT OF HEART  
DISEASES IN LOCAL POPULATION OF PAKISTAN**<sup>1</sup>Dr. Muhammad Wajid, <sup>1</sup>Dr. Sidra Afzal, <sup>1</sup>Dr. Ahsan Wali<sup>1</sup>Bahawal Victoria Hospital, Bahawalpur**Abstract:**

**Introduction:** Hypertension is a modifiable and major risk factor for coronary artery disease, heart failure, cerebrovascular disease and chronic renal failure. It is also recognized as a global chronic, non-communicable disease and a “silent killer” due to its high mortality rates and lack of early symptoms. **Aims and objectives:** The main objective of the relationship of blood pressure and hypertension in the risk assessment of heart diseases in local population of Pakistan. **Methodology of the study:** This study was conducted at Bahawal Victoria hospital, Pakistan during 2018. The data was collected from 100 patients which was suffering from high blood pressure and any kind of heart issue. We collect the data in two sections, as first of all we collect some demographic information regarding age, sex, socio-economic status and history of blood pressure. Then in second part we collect data regarding high blood pressure and heart issues. For this purpose we prepare a questionnaire and fill that from patients. **Results:** The data shows that there is a significant relationship between hypertension and CVD. There is also some positive relationship between socio-economic status and hypertension with respect to CVD. Table 01 shows the value of LDL, HDL, Cholesterol and demographic values of patients. **Conclusion:** In conclusion, the current hypertension paradigm does not account for the continuous risk associated with elevated BP or the multifactorial nature of CVD, the primary consequence of elevated BP.

**Corresponding author:**

**Dr. Muhammad Wajid,**  
Bahawal Victoria Hospital,  
Bahawalpur

QR code



Please cite this article in press Muhammad Wajid et al., *A Study on Relationship of Blood Pressure and Hypertension in the Risk Assessment of Heart Diseases in Local Population of Pakistan.*, Indo Am. J. P. Sci, 2018; 05(11).

**INTRODUCTION:**

Hypertension is a modifiable and major risk factor for coronary artery disease, heart failure, cerebrovascular disease and chronic renal failure. It is also recognized as a global chronic, non-communicable disease and a “silent killer” due to its high mortality rates and lack of early symptoms. One-quarter of the world’s adult population is hypertensive and it is estimated that by 2025 this figure is likely to increase to 29% [1]. These relations are steady in the two people, in youthful, moderately aged, and more seasoned subjects, among different racial and ethnic gatherings, and inside and between nations. In spite of the fact that there is a continuum of cardiovascular hazard crosswise over levels of circulatory strain, the characterization of grown-ups as indicated by pulse gives a system to differentiating levels of hazard related with different circulatory strain classes and for characterizing treatment edges and helpful objectives [2].

As per the grouping approaches created by the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC VI) and the World Health Organization and the International Society of Hypertension (WHO-ISH), non-hypertensive subjects with a systolic weight of 130 to 139 mm Hg or a diastolic weight of 85 to 89 mm Hg are sorted as having high-ordinary pulse [4]. Despite the fact that subjects with high-ordinary circulatory strain are probably going to have a hoisted danger of cardiovascular infection (given the continuum of hazard), there is a scarcity of data in regards to the supreme and relative dangers of cardiovascular ailment in these people [5]. In spite of the fact that information on deadly coronary occasions and strokes in people with high-typical circulatory strain are accessible, data on the danger of nonfatal cardiovascular occasions among individuals in this pulse class is restricted. We attempted a planned examination of the danger of cardiovascular sickness in people with high-typical pulse [6].

Elevated blood pressure (BP) is a causal risk factor for cardiovascular disease (CVD). In addition,

randomized clinical trials among people with hypertension have illustrated, in total, a decrease in CVD occasions by 20%, coronary illness (CHD) by 17%, stroke by 27%, and heart disappointment by 28% for each 10 mm Hg systolic BP (SBP) bringing down with medicinal treatment. In this manner, counteractive action, location, treatment, and control of lifted BP, and its clinical connect hypertension, is a critical general health need and an essential focus for CVD aversion [7].

**Aims and objectives**

The main objective of the relationship of blood pressure and hypertension in the risk assessment of heart diseases in local population of Pakistan.

**METHODOLOGY OF THE STUDY:**

This study was conducted at Bahawal Victoria hospital, Pakistan during 2018. The data was collected from 100 patients which was suffering from high blood pressure and any kind of heart issue. We collect the data in two sections, as first of all we collect some demographic information regarding age, sex, socio-economic status and history of blood pressure. Then in second part we collect data regarding high blood pressure and heart issues. For this purpose we prepare a questionnaire and fill that from patients.

**Statistical analysis**

Student’s t-test was performed to evaluate the differences in roughness between group P and S. Two-way ANOVA was performed to study the contributions. A chi-square test was used to examine the difference in the distribution of the fracture modes (SPSS 19.0 for Windows, SPSS Inc., USA).

**RESULTS:**

The data shows that there is a significant relationship between hypertension and CVD. There is also some positive relationship between socio-economic status and hypertension with respect to CVD. Table 01 shows the value of LDL, HDL, Cholesterol and demographic values of patients.

**Table 01:** Statistical analysis values of Control group and diseased group

Variable	Diseases Group	Control Group	t Value	p Value
Age (Year)	56.56±8.46	53.64±8.36	1.716	0.081
BMI (kg/m <sup>2</sup> )	24.31±2.26	23.37±2.09	2.195	0.031
SBP (mmHg)	140.36±15.70	116.53±13.46	8.248	0.000
DBP (mmHg)	87.94±10.69	75.81±9.94	5.967	0.000
PP (mmHg)	52.42±12.87	40.72±8.74	5.426	0.000
FBG (mmol/)	5.12±0.65	5.06±0.49	1.764	0.081
TG (mmol/L)	1.74±0.75	1.69±0.86	1.838	0.071
TC (mmol/L)	4.95±0.76	4.88±0.82	1.712	0.090
HDL-	1.30±0.43	1.31±0.56	1.717	0.089
LDL-C	3.46±0.58	3.38±0.66	1.139	0.266

**DISCUSSION:**

Our approach to understand disease development in early life, identify key pathways of interest in predisposition to hypertension and develop specific preventive approaches has been to use multi-modality imaging to capture information on cardiovascular structure and function ‘from heart to capillary’. With this approach it becomes possible to model the interrelationship between features of the cardiovascular system and, with longitudinal data, study the progression of disease across vessel and heart. By extending the data collection to other organs such as brain and liver, a holistic view of disease development can be captured [8].

High blood pressure was the leading risk factor for the overall global burden of disease in 2010. The recent decrease in cardiovascular mortality in high-income countries has been associated with a rise in the numbers of patients living with cardiovascular disease, and the wider use of preventive drugs [9]. Thus, an up-to-date understanding of the associations of blood pressure with different non-fatal and fatal cardiovascular disease outcomes would help to refine strategies for primary prevention and inform the design of future clinical trials [10].

The National Health Survey of Pakistan estimated that hypertension affects 18% of adults and 33% of adults above 45 years old. In another report, it was shown that 18% of people in Pakistan suffer from hypertension with every third person over the age of 40 becoming increasingly vulnerable to a wide range of diseases. It was also mentioned that only 50% of the people with hypertension were diagnosed and that only half of those diagnosed were ever treated. Thus, only 12.5% of hypertension cases were adequately

controlled. Some remote areas like Balochistan, there is a paucity of data but the control rate is likely to get even worse [11].

**CONCLUSION:**

In conclusion, the current hypertension paradigm does not account for the continuous risk associated with elevated BP or the multifactorial nature of CVD, the primary consequence of elevated BP.

**REFERENCES:**

1. Lim SS, Vos T, Flaxman AD. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224–2260
2. Hippisley-Cox J, Coupland C, Robson J, Brindle P. Derivation, validation, and evaluation of a new QRISK model to estimate lifetime risk of cardiovascular disease: cohort study using QRisk database. *BMJ*. 2010;341:c6624.
3. Lloyd-Jones DM, Leip EP, Larson MG, Vasan RS, Levy D. Novel approach to examining first cardiovascular events after hypertension onset. *Hypertension*. 2005;45:39–45.
4. Herrett E, Shah AD, Boggon R. Completeness and diagnostic validity of recording acute myocardial infarction events in primary care, hospital care, disease registry, and national mortality records: cohort study. *BMJ*. 2013;346:f2350.
5. Gallagher AM, Puri S, van Staa TP. Linkage of the General Practice Research Database (GPRD) with other data sources. *Pharmacoepidemiol Drug Saf*. 2011;20:S230–S367.

6. Beckett N, Peters R, Tuomilehto J, the HYVET Study Group Immediate and late benefits of treating very elderly people with hypertension: results from active treatment extension to hypertension in the very elderly randomised controlled trial. *BMJ*. 2012;344:d7541.
7. Murabito JM, Evans JC, Nieto K, Larson MG, Levy D, Wilson PW. Prevalence and clinical correlates of peripheral arterial disease in the Framingham Offspring Study. *Am Heart J*. 2002;143:961–965.
8. Goff DC, Jr, Lloyd-Jones DM, Bennett G. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*.
9. JBS3 Board Joint British Societies' consensus recommendations for the prevention of cardiovascular disease (JBS3) *Heart*. 2014;100(suppl 2):ii1–i67.
10. Selvin E, Erlinger TP. Prevalence of and risk factors for peripheral arterial disease in the United States: results from the National Health and Nutrition Examination Survey, 1999–2000. *Circulation*. 2004;110:738–743
11. Azhar S, Hassali MA, Ibrahim MI, et al. The role of pharmacists in developing countries: the current scenario in Pakistan. *Hum Res Health*. 2009;7:54.