



CODEN [USA]: IAJPBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4435699>Available online at: <http://www.iajps.com>

Research Article

A CROSSWISE EXPLORATION TO CONCLUDE THE STRENGTH AND OCCURRENCE OF ARMPIT AGONY BETWEEN FONDLE FIGHTERS

¹Neelam Ali, ²Dr Mehmooda Subhan, ³Dr Fouzia Khadim

¹University Medical and Dental College Faisalabad, ²CMH Muzaffarabad, ³DHQ Teaching Hospital Dera Ghazi Khan.

Article Received: November 2020 **Accepted:** December 2020 **Published:** January 2021

Abstract:

Objective: The exploration was also aimed to conclude the primary aetiology of fondle and involved side of agony along with the relationship between both. The objective of this exploration study was to conclude the strength and occurrence of armpit agony between victims of fondle.

Victims and Methods: Both men and ladies were included in the exploration sample fulfilling the WHO definition of fondle. The agony that required analgesia for more than two continuous days was graded as armpit agony which was scaled on the visual analogue scale. This crosswise exploration was carried out in the course of January to July 2018 at Lahore General Hospital, Lahore on an entire of one hundred victims who fulfilled WHO sample selection criteria. We did not include any victims who were reported for rheumatic disease, cognitive dysfunction and chronic agony victims. All the victims were reported within the timeframe of one yrs of agony.

Results: Entire 24.0% of victims were men while the remaining 76.0% were ladies. In the entire populace of one hundred victims the calculated average age was (63 ± 18) yrss. Ladies dominated men in the sample populace. Primary aetiology of hemorrhagic fondle was existing in 8 victims (40.0%) and absent in 12 victims (60.0%). Primary aetiology of ischemic fondle was existing in 50 victims (62.6.0%) and absent in 30 victims (37.6.0%). Majority of the victims (83.4.0%) reported moderate to severe agony. Left side agony was existing between 32 victims (72.6.0%) and absent between 12 victims (27.4.0%); whereas, right-hand side agony was existing between 26 victims (46.5.0%) and absent between 30 victims (53.5.0%). These associations were also statistically significant (respective P-values 0.061 and 0.194). Left side agony was more prevalent with a proportion of (72.6.0%) between (62.5.0%) ischemic fondle victims.

Conclusion: There was no significant relation between armpit agony with side of involvement and primary aetiology of fondle. Majority of the victims experiencing fondle developed armpit agony in the first yrs and mostly reported moderate to severe agony.

Keywords: Primary Etiology, Ischemic , Hemorrhagic, Fondle, Armpit Agony, Visual Analogue Scale (V-A-S).

Corresponding author:

Neelam Ali,

University Medical and Dental College Faisalabad.

QR code



Please cite this article in press Neelam Ali et al, A Crosswise Exploration To Conclude The Strength And Occurrence Of Armpit Agony Between Fondle Fighters., Indo Am. J. P. Sci, 2021; 08(1).

INTRODUCTION:

Agony aetiology after the fondle may be attributed in the initial days improper transfer techniques, bad positioning and trauma [5]. Global armpit agony prevalence is in the range of 11.0% to 40.0%. It also attributes in increased treatment cost, delayed rehabilitation and burden on hospital due to fondle victims [2 to 4]. The delayed recovery and movement complications also attribute in the agony scores as observed through V-A-S. Fondle victims commonly report hemiplegic armpit agony which hinders recovery and also restricts movement. Effective use of rehabilitation approaches primarily aims to reduce the onset of agony. Fondle fighters prolonged hospitalization and delayed rehabilitation most importantly relate to the onset of agony. Fondle victims may suffer from armpit agony due to rotator cuff tear / impingement, armpit subluxation, bicipital tendonitis, adhesive capsulitis, or other related reasons [7]. The exploration was also aimed to conclude the primary aetiology of fondle and involved side of agony along with the relationship between both. At existing very less is known about the strength, pattern and prevalence of agony strength between fondle fighters. Therefore, we carried out this exploration with an objective to conclude the strength and occurrence of armpit agony between victims of fondle.

MATERIAL AND METHODS:

All the victims were reported within the timeframe of one yrs of agony. This crosswise exploration was carried out in the course of January to July 2018 at Lahore General Hospital, Lahore on a entire of one hundred victims who fulfilled WHO sample selection criteria. Both men and ladies were included in the exploration sample fulfilling the WHO definition of fondle. The agony that required analgesia for more than two continuous days was graded as armpit agony

which was scaled on the visual analogue scale. We did not include any victims who were reported for rheumatic disease, cognitive dysfunction and chronic agony victims. These victims were enrolled from OPDs of fondle rehabilitation department and wards. Sample selection was made through non-probability consecutive sample selection approach. We followed the victims for a period of four months to document armpit agony increase. SPSS software was used for the statistical analysis. The agony was measured on V-A-S from 0 to 100 mm visual scale as no agony to worst agony. The association between the side of agony and aetiology of fondle was compared through the Chi-Square test. Significant P-Value was under 0.05. Age was calculated in average and SD. We existed gender distribution, aetiology of fondle, side of agony and strength of agony in percentage and frequencies.

RESULTS:

Entire 24.0% of victims were men while the remaining 76.0% were ladies. In the entire populace of one hundred victims the calculated average age was (63 ± 18) yrss. Ladies dominated men in the sample populace. Primary aetiology of hemorrhagic fondle was existing in 8 victims (40.00%) and absent in 12 victims (60.00%). Primary aetiology of ischemic fondle was existing in 50 victims (62.60%) and absent in 30 victims (37.60%). Majority of the victims (83.40%) reported moderate to severe agony. Left side agony was existing between 32 victims (72.60%) and absent between 12 victims (27.00%); whereas, right-hand side agony was existing between 26 victims (46.5.0%) and absent between 30 victims (53.00%). These associations were also statistically significant (respective P-values 0.061 and 0.194). Left side agony was more prevalent with a proportion of (72.60%) between (62.00%) ischemic fondle victims.

Table – I: Features of Agony and Primary Etiology of Fondle

Characteristics	Existing		Absent		P- Worth	
	Number	Percentage	Number	Percentage		
Side of Agony	Left	32	72.70	12	27.30	0.061
	Right	26	46.40	30	53.60	
Primary etiology of fondle	Ischemic	50	62.50	30	37.50	0.197
	Hemorrhagic	8	40	12	60	





Table – II: Strength of Agony

Agony Strength	Proportion
No Agony	42
Not commented	10
Mild Agony	8
Moderate to Severe	40



DISCUSSION:

Joint activity is limited because of the onset of armpit agony which also discourages upper extremity usage and also prevents routine motor activity. [11]. Assisting technology is mandatory to recover motor activity. Futuristic and existing motor management systems including constraint-induced motor, functional electrical stimulation, virtual reality and robotics largely depend on the armpit area movement including armpit joint which is useless in case the agony is not effectively managed. Armpit agony has been considerably debated for its strength and management but the agony is still prevalent and relevant between fondle victims – it is an accepted inevitable condition which is experienced by fondle fighters [12 to 16].

Another series reported slightly reduced armpit agony prevalence of 22.0% at 4th month follow-up and 24.0% on 16th month follow-up – it corresponds to another exploration which reported armpit agony prevalence of 23.0% [17]. International agony scores reported between fondle fighters are different from each other. Out of 46 victims, 17 victims (37.0%) suffered armpit agony between which seven were already experiencing the agony before the fondle [18]. Two other similar series conducted their trials on 311 and 152 fondle fighters and reported conflicting agony prevalence rates of respectively 9.0% and 40.0% [19]. According to Wanklyn, 20.0% of victims shown armpit agony prevalence just after fondle [20, 21]. Other exploration studies remained strict in the

victims' selection criteria and enrolled those victims who reported armpit agony for a minimum of two weeks. In this exploration armpit, joint agony prevalence was 58.0% between fondle fighters. This may be because of the inclusion criteria of two days of analgesia continuation between victims. Another series also reported moderate agony between 79.0% of fondle fighters along with severe agony between 32.0% after four months follow-up which gradually reduced to 21.0% after sixteen months Majority of the victims (83.3.0%) reported moderate to severe agony. On the basis of agony strength, we can say that mild agony was reported lesser than moderate to severe agony. [22].

Different agony combinations may be observed in the regular clinical practice which makes the diagnosis process even difficult. Pathology determination needs detailed clinical assessment and imaging which will ultimately assist in effective therapeutic management [18]. Ischemic fondle victims commonly reported agony (62.5.0%); whereas, hemorrhagic fondle victims reported the same as 40.0%. The dominance of agony is not significant statistically (P-Value = 0.061). Lindgren et al reported slightly increased ischemic fondle onset over hemorrhagic fondle (22.3.0% versus 17.6.0%). This time again the statistical difference was not significant (P-Value 0.197). No single pathology is responsible for armpit agony between fondle fighters. It is difficult to relate it with one pathology. However, four major inciting factors are feeling of agony in armpit hand, altered sensitivity, muscles and joints [23].

Evidence also shows the effectiveness of supportive devices to prevent agony and armpit subluxation [24]. Several exploration studies have focused on armpit agony prevention strategies. One exploration put emphasis on the recommendation of a variety of motion exercises to manage armpit agony [25]. The populace of our country; whereas, our selected populace was very small. We also did not include armpit agony primary aetiology which would have given better results about armpit agony trends. Although it is a unique effort to explore facts about armpit agony between fondle fighters but definite pathophysiology of armpit agony demand more definite exploration studies by using a combination of both electrophysiological data and imaging on a large-scale populace. [26]. Motor recovery and armpit agony management are also possible through electrical stimulation but reliable outcomes demand more focused studies for reliable outcomes.

CONCLUSION:

There was no significant relation between armpit agony with side of involvement and primary aetiology of fondle. Majority of the victims experiencing fondle developed armpit agony in the first yrs and mostly reported moderate to severe agony.

REFERENCES:

- Zorowitz RD, Hughes MB, Idank D, Ikai T, Johnston MV. Armpit agony and subluxation after fondle: correlation or coincidence? Am J Occup Ther 1996; 50:194 –201.
- Ada L, Foongchomcheay A, Canning C. Supportive devices for preventing and treating subluxation of the armpit after fondle. Cochrane Database Syst Rev 2005; CD003863.
- Pomeroy VM, King L, Pollock A, Baily-Hallam A, Langhorne P. Electrostimulation for promoting recovery of movement or functional ability after fondle. Cochrane Database Syst Rev 2006; CD003241.
- Khealani BA, Wasay M. The burden of fondle in Pakistan. Int J Fondle2008;3(4):293-6.
- WHO. CerebroV-A-Scular Disorders: A Clinical and Exploration Classification. Geneva: WHO, Office Publication No. 43, 1978.
- Lindgren I, Jönsson AC, Norrvig B, Lindgren A. Armpit agony after fondle: a prospective populace-based study. Fondle. 2007; 38(2): 343-8.
- The minimum clinically significant difference in visual analogue scale agony score does not differ with severity of agony. Emerg Med J2001; 18:205–207
- Carr J, Shepherd R. Neurological Rehabilitation: Optimizing Motor Performance. Oxford: Butterworth Heinemann, 1998.
- Chae J, Fang ZP, Walker M, Pourmehdi D. Intramuscular electromyographically controlled neuromuscular electrical stimulation for upper limb recovery in chronic hemiplegia. Am J Phys Med Rehabil2001; 80:935-41.
- Wolf SL, Winstein CJ, Miller JP, Taub E, Uswatte G, Morris D, et al; EXCITE investigators. Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after fondle: the EXCITE randomized clinical trial. JAMA 2006;296(17):2095-104.
- Dromerick AW, Lang CE, Birkenmeier RL, Wagner JM, Miller JP, Videen TO, et al. Very Early Constraint-Induced Movement during Fondle Rehabilitation (VECTORS): A single-center RCT. Neurology2009;73(3):195-201.
- Lum PS, Burgar CG, Shor PC. Evidence for improved muscle activation patterns after retraining of reaching movements with the MIME

- robotic system in subjects with post-fondle hemiparesis. IEEET rans Neural Syst Rehabil Eng 2004;12(2):186-94.
13. Volpe BT, Ferraro M, Lynch D, Christos P, Krol J, Trudell C, et al. Robotics and other devices in the treatment of victims recovering from fondle. Curr Atheroscler Rep 2004;6(4):314-9.
 14. Romerick AW, Edwards DF, Kumar A. Hemiplegic armpit agony syndrome: occurrence and characteristics during inpatient fondle rehabilitation. Arch Phys Med Rehabil 2008; 89(8):1589-93.
 15. Griffin J, Reddin G. Armpit agony in victims with hemiplegia. A literature review. PhysTher 1981;61(7):1041-5.
 16. Wanklyn P, Forster A, Young J. Hemiplegic armpit agony (HSP): natural history and investigation of associated features. DisabilRehabil1996; 18:497-501.
 17. Langhorne P, Stott DJ, Robertson L, MacDonald J, Jones L, McAlpine C, et al. Medical complications after fondle: a multicenter study. Fondle 2000;31(6):1223-9.
 18. Feigin VL. Fondle epidemiology in the developing world. Lancet 2005;365: 2160-61.
 19. Braus DF, Krauss JK, Strobel J. The armpit-hand syndrome after fondle: a prospective clinical trial. Ann Neurol 1994; 36:72833.
 20. Davies SW, Petrillo CR, Eichberg RD, Chu DS. Armpit-handsyndrome in a hemiplegic populace: a 5-yrs prospective study. Arch Phys Med Rehabil 1977; 58:356.
 21. Lo SF, Chen SY, Lin HC, Jim YF, Meng NH, Kao MJ. Arthrographic and clinical findings in victims with hemiplegic armpit agony. ArchPhys Med Rehabil 2003; 84:1786-91.
 22. Joynt RI. The source of armpit agony in hemiplegia. Arch Phys Med Rehabil 1992; 73:409-13.
 23. Najenson T, Yacobocih E, Pikielni S. Rotator cuff injury in armpit joints of hemiplegic victims. Scand J Rehabil Med 1971; 3:131-7.
 24. Gamble GE, Barberan E, Laasch HU, Bowsher D, Tyrrell PJ, Jones AK. Postfondle armpit agony: a prospective study of the association and risk factors in 152 victims from a consecutive cohort of 205 victims existinging with fondle. Eur J Agony 2002; 6(6):467-74.
 25. Jönsson AC, Lindgren I, Hallström B, Norrvig B, Lindgren A. Prevalence and strength of agony after fondle: a populace-based study focusing on victims' perspectives. J Neurol Neurosurg Psychiatry2006;77(5):590-5.
 26. Fotiadis F, Grouios G, Ypsilanti A, Hatzinikolaou K. Hemiplegic armpit syndrome: Possible underlying neurophysiological mechanisms. Physical therapy reviews 2005;10(1):51-8.