



CODEN [USA]: IAJPBB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4445654>Available online at: <http://www.iajps.com>

Research Article

**FIND OUT THE AMALGAMATED AND COMPARATIVE
SYNERGETIC EFFECTS OF METFORMIN AND BLACK
COFFEE IN TYPE 2 DIABETES MELLITUS TREATMENT**Dr. Shanza Kanwal¹, Dr Hafiza Zara Amin², Dr. Farman Ali³¹Ameer-ud-din Medical College, Lahore, ²Sir Ganga Ram Hospital, Lahore, ³Gomal Medical College, Dera Ismail Khan.

Article Received: November 2020 Accepted: December 2020 Published: January 2021

Abstract:

Objective: The aim of our study was to find out the amalgamated and comparative synergetic effects of metformin and black coffee in type 2 diabetes mellitus treatment.

Study Design: An experimental, randomized control study.

Place and Duration: This study was conducted at Pharmacology laboratory and Multidisciplinary laboratory of Services hospital Lahore for the duration of one year starting from November, 2019 to October, 2020.

Methodology: In our study we enrolled 50 male weighing 30-50 g albino Balb/c mice. Group 1 was consisting on non-diabetic normal control (n=10) and diabetes was induced in experimental group (n=40) by using low dose streptozotocin (40mg/kg). We further divide diabetic mice into 4 groups (10 mice/group). Group 2 consist on diabetic controls and the remaining three groups treated with metformin, black coffee and combination of both for 45 days. We use intracardiac puncture method to took blood samples for HbA1c. SPSS version 20 were used for the statistical analysis of data. With the use of one-way ANOVA (post hoc Tuckey test), comparisons of HbA1c among different groups were analyzed. (P-value <0.05)

Results: According to the results of our study we noticed that metformin treated (Group 4), black coffee treated (Group 3) and combination of black coffee and metformin treated (Group 5) had decreased the level of serum HbA1c in comparison with those who found in diabetic control (Group 2). (P<0.05)

Conclusion: At the end of our study, we conclude that combination of metformin and black coffee decrease the level of serum of HbA1c in diabetic mice as compared to separately use of black coffee and metformin.

Key Words: HbA1c, Diabetes Mellitus, Pancreatic Islets, Cells Metformin

Corresponding author:**Dr. Shanza Kanwal**

Ameer-ud-din Medical College, Lahore.

QR code



Please cite this article in press Shanza Kanwal et al, Find Out The Amalgamated And Comparative Synergetic Effects Of Metformin And Black Coffee In Type 2 Diabetes Mellitus Treatment., Indo Am. J. P. Sci, 2021; 08[1]

INTRODUCTION

Diabetes mellitus is spreading worldwide pandemic that is a raising wellbeing concern with numerous difficulties and an expanding ubiquity [1]. Despite stunning change over both essential and clinical therapeutic sciences, diabetes mellitus is at present a hopeless life-long sickness, rapidly affecting both genders [2]. Diabetes mellitus may be not a single illness rather an aggregation of metabolic issue connected with secondary harm on various organ framework that incorporate cardiovascular diseases, stroke, nephropathy, retinopathy, neuropathy, gangrene and even amputations [3,4]. Diabetes mellitus affects 382 million people globally [5,6]. The current prevalence of this disease in Pakistan is 11.77% according to a study conducted in 2015 [7]. The pandemic of type 2 diabetes mellitus has been met by emerging approach and clinical tactics, including the generally-accepted commendation to institute drug therapy concomitant with lifestyle changes [8,9]. Metformin an older and broadly acknowledged prime agent, has antihyperglycemic properties and also other important functions such as enhancement in endothelial dysfunction, hemostasis and oxidative stress, insulin resistance, lipid profiles, and fat redistribution [10]. Metformin's efficacy, security profile, beneficial cardiovascular and metabolic effects and its capacity to be associated with other antidiabetic agents makes this drug the first glucose lowering agent of choice when treating patients with type 2 diabetes [11,12].

Even the most explored out oral antidiabetic drugs sometimes fail as monotherapy and eventually different drug combinations are to be considered [13]. Many new trends in management of diabetes are therefore being considered nowadays and the approach for diabetes treatment is moving towards the incorporation of more organic natural products which tend to counter the root causes of the disease [14,15]. Black coffee is the most commonly used energy boosting beverage worldwide. Recent studies have shown many potential health benefits of black coffee in humans [16]. Besides keeping you alert and wakeful, coffee is the richest and intense source of antioxidants that work as meager warriors battling and securing against free radicals inside human body [17]. Caffeine in black coffee regulates the hyperglycemic effect in diabetic patients by increasing insulin release from pancreatic beta cells by the sensitization of the ryanodine receptor and activation of 5'-adenosine monophosphate-activated protein kinase (AMPK). Caffeine also up regulates the insulin-like growth factor 1 signaling, which is responsible for enhanced insulin sensitivity as well as insulin secretion. Chlorogenic acid has an eloquent role in glucose

metabolism by decreasing glucose output in the liver and promoting the synthesis of the "homeodomain transcription factor IDX-1", which directs beta cells to counter the increased glucose levels in plasma. Important antioxidants in coffee include hydro cinnamic acids and polyphenols, Hydro cinnamic acids are very effective at neutralizing free radicals and preventing oxidative stress. Polyphenols counter the increased insulin resistance and escalates the insulin sensitivity [18,19,20,21].

The cumulative body of suggestion about lower frequency of diabetes among coffee users is conclusive enough to prove a positive impact of coffee consumption on the development of type 2 diabetes mellitus. Different studies have been conducted to evaluate the preventive role of black coffee on type 2 diabetes [22]. To the best of our knowledge, no study has been conducted to explore the effect of black coffee as adjunct to metformin in treatment of diabetes patients. If this agent works to improve insulin sensitivity and decreases the insulin resistance, then this cost effective and easy administered agent with overall beneficial effects on health can be used in place of other antidiabetic agents who are usually given with metformin, when metformin monotherapy fails. This experimental study was done to determine the comparative and combined synergistic effect of Black Coffee and Metformin in treatment of type 2 diabetes mellitus in mice model.

METHODOLOGY:

This randomized control study was conducted at Pharmacology laboratory and Multidisciplinary laboratory of Services hospital Lahore for the duration of one year starting from November, 2019 to October, 2020. A total of 50 healthy, 6-8 weeks old male, weighing 30-50 g albino Balb/c mice were included in the study. All the mice were accommodated in standard cages which were made up of plastic and placed on metallic racks, at the Animal house of NIH, Lahore. Room number 13 was allocated for the research procedure. The mice had free access to tap water through the inverted bottles of 250ml capacity fixed on top of the cages. These bottles were cleaned and filled on daily basis according to the protocol of the animal house. The normal standard diet was prepared at the NIH, which was served with standard food pellets. Animal house atmosphere was maintained at room temperature of 20 ± 2 oC with relative humidity of 50-70% with a light and dark cycle of 12 hours each. After acclimatization for 1 week, the mice were randomly divided into two groups; 10 mice were allocated to Group 1 and

remaining 40 mice were allocated to the Experimental Group.

Group 1 was labeled as Normal Control and was given normal diet for 5 days whereas the Experimental group was given normal diet plus streptozotocin, (STZ), (40mg /kg /day) 2 3 intraperitoneally for consecutive 5 days. After 5 days, confirmation of diabetes in experimental group was done by measuring and comparing fasting blood glucose levels (mg/dl) with Group 1. The blood sample was taken from lateral tail vein of all mice with 1 ml syringe and the blood glucose levels were measured by using EASY GLUCO Ultra Plus Auto Coding meter Iso tech Co. Ltd. Experimental group was then further divided into four groups i.e., 2 (Diabetic control), 3 (Black coffee treated), 4 (Metformin treated) and 5(Combination of Black coffee and Metformin treated).

Group 2 mice were given normal standard diet only. Group 3 mice were given normal diet mixed with Black Coffee (5g/kg/day)²⁴ orally for 45 days. Group 4 mice were given normal diet along with Metformin (200mg/kg/day)²⁵ orally mixed in drinking water for 45 days. Group 5 mice were given normal diet mixed with Black Coffee (5g/kg/day) orally and Metformin

(200mg/kg/day) orally mixed in drinking water for 45 days. After 45 days of treatment, final sampling of the experiment was done from group 3, 4 and 5 which included HbA1C (%) by cardiac puncture. Fixed time nephelometry certified by National Glycohemoglobin Standardization Program (NGSP) was employed for HbA1C estimation (%). For this study, PA50 fully auto specific protein analyzer was used. Statistical analysis was done by applying the Statistical Package for Social Sciences version 21 (SPSS 21). Results were documented as Mean \pm Standard Error of Mean (SEM). Comparison of means of HbA1C (%) among the five groups were analyzed by using the One-way ANOVA and Post hoc Tuckey tests. P value of <0.05 was considered significant.

RESULTS:



The result of Mean \pm SEM of HbA1C (%) in group 2 (7.82 \pm 0.11) was significantly higher than group 1 (P value >0.05). While on comparison of Mean \pm SEM of HbA1C (%) in group 3 (6.02 \pm 0.29), group 4 (5.76 \pm 0.45) and group 5 (4.90 \pm 0.28) were significantly lower than group 2 (P value <0.05). Table 01 shows the comparison of mean \pm SEM of HbA1C (%) of all the groups.

Table No 01: Comparison of Mean \pm SEM of HbA1C (%) in all five Groups

Groups	Mean \pm SEM	P-value
Control	4.75 \pm 0.13	<0.05
Diabetes	7.82 \pm 0.11	
Black Coffee	6.02 \pm 0.29	
Metformin	5.76 \pm 0.45	
Combination of Black Coffee and Metformin	4.90 \pm 0.28	

Table No 02: Multiple Comparison of Mean Difference of Hba1c (%) of Control and Experimental Groups


Groups	Mean Difference (%)	P-value
Group 1 vs group 2	3.07%	0.000*
Group 1 vs group 3	1.27%	0.020
Group 1 vs group 4	1.01%	0.102
Group 1 vs group 5	0.15%	0.995
Group 2 vs group 3	1.79%	0.000*
Group 2 vs group 4	2.06%	
Group 2 vs group 5	2.91%	
Group 3 vs group 4	0.26%	0.961
Group 3 vs group 5	1.12%	0.053
Group 4 vs group 5	0.85%	0.219



The graphical representation of HbA1C (%) results shows a marked difference between the Black coffee, Metformin, Combination of Black coffee and Metformin as compared to the diabetic control mice. Metformin has a better role in lowering HbA1C (%) than black coffee, yet the role of combination therapy is astonishing in aspect to lower HbA1C (%) nearly to the normal control range.

Table No 03: HbA1C (%) Results in Different Groups

Groups	HbA1C (%)
Control	5.00
Diabetes	7.85
Black Coffee	6.00
Metformin	5.65
Combination	5.23



DISCUSSION:

The results of present study confirm that hyperglycemia induced by streptozotocin, is ameliorated by all the experimental agents to an appreciable extent, yet the result of combination therapy of Black coffee and Metformin is very impressive. In present study, the antidiabetic effect of Metformin is seen in group 4 and in combination with Black coffee in group 5. Improvement of HbA1C in Group 4 is supported by study of S.H Chung *et al.*, who compared the antidiabetic effect of metformin and compound k in diabetic db/db mice and proposed that normalization of raised plasma glucose levels and improvement in insulin levels in metformin treated group [26]. In present study, improvement in HbA1C in Groups 3 and 5 along with better improvement in Group 5 which is given combination of black coffee and metformin is observed.

Kobayashi M *et al.*, also demonstrated similar results who used black coffee, caffeine extract, decaffeinated coffee against different sets of experiments to analyze the preventive part of black coffee on development of STZ induced diabetes and also the reversal of worsening offered by STZ induced hyperglycemia in male C57 BL/ 6J mice. He demonstrated that continuous Black coffee ingestion prevented the development of STZ induced diabetes mellitus and also revealed that the black coffee can recover the hyperglycemia induced metabolic changes by analyzing the biochemical and histopathological parameters [27]. Mukesh Doble *et al.*, did a study to demonstrate comparative and combined effects of plant phenolic compounds, chlorogenic acid and ferulic acid with metformin and thiazolidinedione on the uptake of 2-deoxyglucose (2DG) by L6 myotubes of rats. He established that a combination of different concentrations of chlorogenic acid and metformin or THZ, has a synergistic effect in the uptake of 2DG with a maximum of 5.0 and 5.3 times respectively, when contrasted to the control. Ferulic acid in combination with metformin or THZ has likewise displayed a synergistic impact and the 2DG uptake increases by 4.98 and 5.11-fold when compared to the control [28]. Hence when Metformin is given in combination with black coffee, HbA1C which shows the long-term control of diabetes is improved demonstrating that black coffee can be used as adjunct to metformin in the treatment of type 2 diabetes mellitus.

CONCLUSIONS:

At the end of our study, we conclude that combination of metformin and black coffee decrease the level of

serum of HbA1c in diabetic mice as compared to separately use of black coffee and metformin.

REFERENCES:

1. Association AD. Diagnosis and classification of diabetes mellitus. *Diabetes care*. 2014;37(Supplement 1): S81-S90.
2. Association AD. Diagnosis and classification of diabetes mellitus. *Diabetes care*. 2012;35(Supplement 1): S64-S71.
3. Nathan DM, Group DER. The diabetes control and complications trial/epidemiology of diabetes interventions and complications study at 30 years: overview. *Diabetes care*. 2014;37(1):9-16.
4. Sayin N, Kara N, Pekel G. Ocular complications of diabetes mellitus. *World journal of diabetes*. 2015;6(1):92.
5. Gagliardino JJ, Atanasov PK, Chan JC, Mbanya JC, Shestakova MV, Leguet-Dinville P, *et al.* Resource use associated with type 2 diabetes in Africa, the Middle East, South Asia, Eurasia and Turkey: results from the International Diabetes Management Practice Study (IDMPS). *BMJ Open Diabetes Research and Care*. 2017;5(1): e000297.
6. Jansson S, Fall K, Brus O, Magnuson A, Wändell P, Östgren CJ, *et al.* Prevalence and incidence of diabetes mellitus: a nationwide population-based pharmaco-epidemiological study in Sweden. *Diabetic medicine*. 2015;32(10):1319-28.
7. Meo SA, Zia I, Bukhari IA, Arain SA. Type 2 diabetes mellitus in Pakistan: Current prevalence and future forecast. *JPMA The Journal of the Pakistan Medical Association*. 2016;66(12):1637-42.
8. Brietzke SA. Oral Antihyperglycemic Treatment Options for Type 2 Diabetes Mellitus. *Medical Clinics*. 99(1):87-106.
9. Association AD. 13. Management of Diabetes in Pregnancy: Standards of Medical Care in Diabetes—2018. *Diabetes care*. 2018;41(Supplement 1): S137-S43.
10. Rojas LBA, Gomes MB. Metformin: an old but still the best treatment for type 2 diabetes. *Diabetology & metabolic syndrome*. 2013;5(1):6.
11. Qaseem A, Barry MJ, Humphrey LL, Forciea M, for the Clinical Guidelines Committee of the American College of P. Oral pharmacologic treatment of type 2 diabetes mellitus: A clinical practice guideline update from the American college of physicians. *Annals of Internal Medicine*. 2017;166(4):279-90.
12. Chakraborty A, Chowdhury S, Bhattacharyya M. Effect of metformin on oxidative stress, nitrate stress and inflammatory biomarkers in type 2

- diabetes patients. *Diabetes research and clinical practice*. 2011;93(1):56-62.
13. Association AD. Standards of medical care in diabetes—2016 abridged for primary care providers. *Clinical diabetes: a publication of the American Diabetes Association*. 2016;34(1):3.
 14. Ghorbani A, Shafiee-Nick R, Rakhshandeh H, Borji A. Antihyperlipidemic effect of a polyherbal mixture in streptozotocin-induced diabetic rats. *Journal of lipids*. 2013;2013.
 15. Hui H, Tang G, Go VLW. Hypoglycemic herbs and their action mechanisms. *Chinese Medicine*. 2015;4(1):11.
 16. Cappelletti S, Daria P, Sani G, Aromatario M. Caffeine: cognitive and physical performance enhancer or psychoactive drug? *Current neuropharmacology*. 2015;13(1):71-88.
 17. Esposito F, Morisco F, Verde V, Ritieni A, Alezio A, Caporaso N, et al. Moderate coffee consumption increases plasma glutathione but not homocysteine in healthy subjects. *Alimentary pharmacology & therapeutics*. 2013;17(4):595601.
 18. Peterson AS. Health Benefits of Coffee. Walter L Aument Family Health Center. 2017;2(4).
 19. Zhou X, Li Y, Shi X, Ma C. An overview on therapeutics attenuating amyloid β level in Alzheimer's disease: Targeting neurotransmission, inflammation, oxidative stress and enhanced cholesterol levels. *American journal of translational research*. 2016;8(2):246.
 20. Hall S, Desbrow B, Anoopkumar Dukie S, Davey AK, Arora D, McDermott C, et al. A review of the bioactivity of coffee, caffeine and key coffee constituents on inflammatory responses linked to depression. *Food Research International*. 2015;76:626-36.
 21. Boia R, Ambrosio AF, Santiago AR. Therapeutic opportunities for caffeine and A2A receptor antagonists in retinal diseases. *Ophthalmic research*. 2016;55(4):212-8.
 22. Carlström M, Larsson SC. Coffee consumption and reduced risk of developing type 2 diabetes: a systematic review with meta-analysis. *Nutrition reviews*. 2018.
 23. Arulmozhi DK, Kurian R, Bodhankar SL, Veeranjanyulu A. Metabolic effects of various antidiabetic and hypolipidemic agents on a high-fat diet and multiple low-dose streptococci (MLDS) mouse model of diabetes. *Journal of Pharmacy and Pharmacology*. 2018;60(9):116773.
 24. Harvei S. Coffee intake and the effects on intestinal inflammation, metabolic homeostasis and intestinal barrier function in mice: Norwegian University of Life Sciences, Ås; 2016.
 25. Aghaalikhani N, Goodarzi MT, Latifi Z, Farimani AR, Fattahi A. Effects of Different Doses of Metformin on Serum Fatty Acid Composition in Type 2 Diabetic Rats. *Avicenna J Med Biochem*. 2017;5(1):22-8.
 26. Wang JY, Zhu C, Qian TW, Guo H, Wang DD, Zhang F, et al. Extracts of black bean peel and pomegranate peel ameliorate oxidative stress-induced hyperglycemia in mice. *Experimental and therapeutic medicine*. 2015;9(1):43-8.
 27. Jin S, Chang C, Zhang L, Liu Y, Huang X, Chen Z. Chlorogenic acid improves late diabetes through adiponectin receptor signaling pathways in db/db mice. *PloS one*. 2015;10(4): e0120842.
 28. Prabhakar PK, Doble M. Synergistic effect of phytochemicals in combination with hypoglycemic drugs on glucose uptake in myotubes. *Phytomedicine*. 2015;16(12):1119-26.