



CODEN [USA]: IAJPBB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

<http://doi.org/10.5281/zenodo.4446346>Available online at: <http://www.iajps.com>

Research Article

**TREATMENT OF TRACHEOESOPHAGEAL FISTULA AND
ESOPHAGEAL ATRESIA; A CLINICAL REVIEW**¹Dr Rana Khurram Aslam, ²Dr Huma Ahmed, ³Dr Nadia Naheed¹University College of Medicine / The University of Lahore, Lahore., ²Rawalpindi Medical University, ³D.G Khan Medical College, Dera Ghazi Khan.**Article Received:** November 2020 **Accepted:** December 2020 **Published:** January 2021**Abstract:**

Aim: To analyze the morbidity and mortality of 38 patients with variants of esophageal atresia and tracheoesophageal fistula treated.

Study design: This is a retrospective study held in the Pediatric Unit-II of Services Hospital Lahore for one-year duration from December 2019 to December 2020. Data was collected from hospital files.

Methodology: The history of 38 patients with esophageal atresia and tracheoesophageal fistula was reviewed.

Results: The mean birth weight was 2.5 kg and the mean gestational age was 38 weeks (range 30 to 38 weeks). The risk classification was based on Waterston's classification and consisted of 18 'A' cases, 12 'B' cases and 8 'C' cases. Many patients have experienced associated abnormalities including heart, skeletal, neurological, and renal and anorectal defects. Anastomotic leakage was found in 8 patients (21.05%), sepsis in 12 (13.5%), and pneumothorax in 3 (7.8%).

Conclusion: The results of this congenital malformation requiring surgical treatment, is less than optimal in our setup. The likely reasons are late presentation, and less than ideal neonatal services.

Keywords: Atresia, fistula, sepsis, congenital anomalies, thoracotomy.

Corresponding author:**Dr. Rana Khurram Aslam,**

University College of Medicine / The University of Lahore, Lahore.

QR code



Please cite this article in press Rana Khurram Aslam et al, *Treatment Of Tracheoesophageal Fistula And Esophageal Atresia; A Clinical Review.*, *Indo Am. J. P. Sci.*, 2021; 08(1).

INTRODUCTION:

The first recorded case of esophageal atresia was in 1670 by Durston in one of the female thoracopagus twins. Thomas Gibsson in 1667 described the most common form of esophageal atresia with a distal fistula. Even after a hundred years of progress in treating esophageal atresia and tracheoesophageal fistula, these newborns continue to be a challenge for pediatric surgeons. The disease incidence is 2.4 per 10,000 live births. Many congenital anomalies are associated with this abnormality. Factors such as low birth weight, delay in diagnosis, late referral, and limited resources play an important role in the poor outcome of our patients. An attempt was made to analyze the factors responsible for morbidity and mortality.

METHODOLOGY:

This is a retrospective study held in the Pediatric Unit-II of Services Hospital Lahore for one-year duration from December 2019 to December 2020. The results of all patients with esophageal atresia and tracheoesophageal fistula were analyzed. Cases were classified into A, B, C, three risk categories based on severity, birth defects, presence or absence of pneumonia, according to Waterston's classification "A" and "B" were operated after initial resuscitation.

The pre-operative tests included basic tests (CBC, blood sugar, urine DR, and serum electrolytes). In all cases, an abdominal ultrasound scan was performed to help recognize any accompanying abdominal deformities. All patients had echocardiography. Right-sided thoracotomy was performed through the 4th intercostal space with an additional pleural access. The lower segment was identified and the fistula ligated and separated from the trachea. The upper pouch was mobilized and a single layer end-to-end anastomosis was made with 4/0 vicryl. A nasogastric tube was passed out of anastomosis and feeding through a nasogastric tube was started on the third day postoperatively. Waterston type "C" patients received complete parental nutrition. Once they were stable, the esophagus and gastrostomy were created and gastrostomy feeding was commenced.

RESULTS:

The mean birth weight was 2.5 kg and the mean gastric age was 38 weeks (range 30 to 38 weeks). According to the Waterston Classification, this group consisted of 18 "A" cases, 12 "B" cases and 8 "C" cases. There were accompanying anomalies in 14 (36.8%) patients, including cardiac and neurological defects, as well as kidney and rectal defects.

TABLE 1: Age at presentation

Age	No of Patients	Percentage
Within 24 Hours	12	31.57
24-48 Hours	10	26.30
More than 48 Hours	16	42.10

Anastomosis leak occurred in 8 (21.05%) patients; sepsis occurred in 12 patients (31.5%). Three patients developed Pneumothorax (7.8%). Late complications included gastroesophageal reflux in 8 (21.05) patients.

TABLE 2: Weight at presentation

Weight	No of Patients	Percentage
Less than 200 gm	4	10.5
> Than 2000 gm	9	23.6
> 2500 gm	10	26.3
More than 3000 gm	15	39.4

Survival in relation to risk categorization is shown in Table 3.

TABLE 3: Western Classification and survival

Category	No. of Patients	Submitted for surgery	Primary staged repair	Survival	Percentage
A	18	14	4	13	72.0
B	12	11	1	06	50.0
C	08		4	01	12.5
Total	38	25	9	20	

25 neonates were treated by primary surgery and repair. Esophagus and inverse gastrostomy were performed in 11 patients, 8 of whom did not have a fistula. The mortality in the study was 47.3% (18 patients)

DISCUSSION:

Esophageal atresia with or without a tracheoesophageal fistula is not a rare birth defect to present at Quetta Children's Hospital, often mimicking feeding difficulties and respiratory distress in the first days of life. The first clinical and pathological description of tracheoesophageal fistula and esophageal atresia was made by Gibbson in 1670. Esophageal atresia with TEF occurs in one of 3,000 to 5,000 births. GPs caring for newborns should be aware of both the clinical symptoms and the management of these newborns. Before the first successful repair was carried out in 1939, the condition was equally disastrous. Over the past 50 years, improvements in neonatal surgical techniques, pre-operative intensive care support, anesthesia and good post-operative care have improved outcomes. An Indian pediatric surgeon reported a large study of 303 cases. Among our patients, 73.68% were men and 26.31% were women, while in the USA the incidence of men is 55%. Table 1 shows that 42% of patients came to us after 48 hours of delivery. This delay in presentation changes the category from "A" to "B" because of pulmonary involvement which adversely affects the prognosis. Over 70% of newborns had a chest infection at admission and required intensive respiratory care. The most common type of esophageal atresia is characterized by a blind proximal esophageal sac with a fistula between the trachea and the distal esophagus. Many anatomical variations of esophageal atresia with and without tracheoesophageal fistula have been described. In this study, 36.81% (14 neonates) had other related defects in addition to esophageal atresia. Rectal anomalies were the most common accompanying anomaly. A wide range of related anomalies, such as ectopic, congenital heart disease, hypospadias. No chromosomal aberrations remained unrecognized due to the inability to conduct the necessary research at the institute. We performed

esophagus and gastrostomy in patients with esophageal atresia with a long gap, with fistula ligation in 4 patients, 3 died. The mortality rate in our patients in this study is 47.3%, mainly due to severe concomitant birth defects and sepsis. In an Italian study, the mortality rate was 92% in 1972, dropped to 55% in 1991 due to a pre-operative evaluation, a good intensive care unit and early surgery. The mortality rate in the large US report is 18%, 6%, and 3% in patients with long intermediate and short interruptions. Spitz et al. In a study published in 1994, reported a range of 3-7.8% mortality according to Waterston's A-C risk categories.

CONCLUSION:

The increase in mortality (47.3%) in this study is mainly due to severe congenital anomalies, sepsis, and delay in diagnosis and referral to pediatric surgery. Survival can be improved with early diagnosis and early referral to a tertiary hospital. The results of this congenital malformation requiring surgical treatment are less than optimal in our system. Late applications and inadequate neonatal services are likely causes.

REFERENCES:

- Schukfeh, Nagoud, Joachim F. Kuebler, Jens Dingemann, and Benno M. Ure. "Thirty years of minimally invasive surgery in children: analysis of meta-analyses." *European Journal of Pediatric Surgery* 30, no. 05 (2020): 420-428.
- Dellenmark-Blom, Michaela, Julia Quitmann, Jens Dingemann, Stefanie Witt, Benno M. Ure, Monika Bullinger, Linus Jönsson, Vladimir Gatzinsky, and Carmen Dingemann. "Clinical factors affecting condition-specific quality-of-life domains in pediatric patients after repair of esophageal atresia: the Swedish-German EA-QOL Study." *European Journal of Pediatric Surgery* 30, no. 01 (2020): 096-103.
- Ax, Sofie Örnö, Kate Abrahamsson, Vladimir Gatzinsky, Linus Jönsson, and Michaela Dellenmark-Blom. "Parent-reported feeding difficulties among children born with esophageal atresia: Prevalence and early risk

- factors." *European Journal of Pediatric Surgery* (2020).
4. Tambucci, Renato, Sara Isoldi, Giulia Angelino, Filippo Torroni, Simona Faraci, Francesca Rea, Erminia Francesca Romeo et al. "Evaluation of Gastroesophageal Reflux Disease 1 Year after Esophageal Atresia Repair: Paradigms Lost from a Single Snapshot?." *The Journal of Pediatrics* 228 (2020): 155-163.
 5. Fraser, Jason D., and Shawn D. St Peter. "The current thoracoscopic management of esophageal atresia." *European Journal of Pediatric Surgery* 30, no. 02 (2020): 142-145.
 6. O'Connor, Elizabeth, and Bruce Jaffray. "Surgeon-Level Variation in Outcome following Esophageal Atresia Repair Is Not Explained by Volume." *European Journal of Pediatric Surgery* (2020).
 7. Bedi, Nandini Kaul, Alka Gupta Grewal, and William Bhatti. "Single institution experience with esophageal atresia over 9 years with a review of literature: Where do we stand?." *CHRISMED Journal of Health and Research* 7, no. 1 (2020): 51.
 8. Koumbourlis, Anastassios C., Yvonne Belessis, Mary Cataletto, Renato Cutrera, Emily DeBoer, Mikhail Kazachkov, Sophie Laberge, Jonathan Popler, Federica Porcaro, and Thomas Kovesi. "Care recommendations for the respiratory complications of esophageal atresia-tracheoesophageal fistula." *Pediatric Pulmonology* 55, no. 10 (2020): 2713-2729.
 9. Oztan, Mustafa O., Tutku Soyer, Can I. Ozturun, Binali Firinci, Çiğdem U. Durakbaşa, Zafer Dokumcu, Gulnur Gollu et al. "Outcome of Very Low and Low Birth Weight Infants with Esophageal Atresia: Results of the Turkish Esophageal Atresia Registry." *European Journal of Pediatric Surgery* (2020).
 10. Dellenmark-Blom, Michaela, Julia Quitmann, and Carmen Dingemann. "Health-related quality of life in patients after repair of esophageal atresia: a review of current literature." *European Journal of Pediatric Surgery* 30, no. 03 (2020): 239-250.
 11. Son, Joonhyuk, Yerang Jang, Wontae Kim, Sanghoon Lee, Ji Seon Jeong, Suk-Koo Lee, and Jeong-Meen Seo. "Thoracoscopic repair of esophageal atresia with distal tracheoesophageal fistula: is it a safe procedure in infants weighing less than 2000 g?." *Surgical Endoscopy* (2020): 1-5.
 12. Elhattab, Ahmad, Liza Ali, Veronique Rousseau, Pauline Clermidi, Daphné Michelet, Caroline Farnoux, Alexandre Lapillonne et al. "Thoracoscopy Versus Thoracotomy in the Repair of Esophageal Atresia with Distal Tracheoesophageal Fistula." *Journal of Laparoendoscopic & Advanced Surgical Techniques* 30, no. 12 (2020): 1289-1294.
 13. Feng, Xiaoyan, Illya Martynov, Anne Suttikus, Martin Lacher, and Steffi Mayer. "Publication Trends and Global Collaborations on Esophageal Atresia Research: A Bibliometric Study." *European Journal of Pediatric Surgery* (2020).
 14. Cui, Xu, Yuanbin He, Liu Chen, Yu Lin, Yunjin Wang, Jianqin Zhang, and Chaoming Zhou. "Clinical Analysis of Azygos Vein Preservation Under Thoracoscope in the Operation of Type III Esophageal Atresia." *Journal of Laparoendoscopic & Advanced Surgical Techniques* 30, no. 4 (2020): 448-452.
 15. Wang, Jiayao, Priyanka R. Ahimaz, Somaye Hashemifar, Julie Khlevner, Joseph A. Picoraro, William Middlesworth, Mahmoud M. Elfiky, Jianwen Que, Yufeng Shen, and Wendy K. Chung. "Novel candidate genes in esophageal atresia/tracheoesophageal fistula identified by exome sequencing." *European Journal of Human Genetics* (2020): 1-9.