



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

SJIF Impact Factor: 7.187

<https://doi.org/10.5281/zenodo.5533367>
Available online at: <http://www.iajps.com>

Research Article

KNOWLEDGE AND PRACTICES OF HEALTHY EATING AND SPORTS AMONG NURSES IN SHEIKH ZAYED HOSPITAL, RAHIM YAR KHAN

¹Parveen Akhtar, ²Naseem Akhtar, ³Tahira Shaheen

¹Charge Nurse, Sheikh Zayed Hospital Rahim Yar Khan, Email ID: parveentayyab94@gmail.com

²Charge Nurse, Sheikh Zayed Hospital Rahim Yar Khan, Email ID:
naseemakhtar13784@gmail.com

³Nursing Instructor, Post Graduate College of Nursing Punjab Lahore Pakistan,
Email ID: ushnamahna@gmail.com

Article Received: July 2021

Accepted: August 2021

Published: September 2021

Abstract:

Introduction: Although nurses are expected to have basic understanding regarding the healthy life style habits and patterns such as balanced diet, physical activity, stress management and hand and sleep hygiene as far as their curriculum is concerned but literature review has shown that they lack insight and urge to translate it into their practical lives for self-grooming and patient education. The purpose of this study is to provide an outlook of the knowledge and practices of the nurses regarding healthy eating and Sports working in Sheikh Zayed Hospital Rahim Yar Khan and formulate recommendations for health-promoting behavior at their workplace. **Methods:** A descriptive study was conducted among 200 nurses working in different departments of Sheikh Zayed Hospital Rahim Yar Khan. A self structured survey questionnaire was used to record the responses from the participants. Data was analyzed by SPSS 21. **Results:** The findings of the study revealed that most of the respondents had heard the term balanced diet early in their lives (82.0%), while (18.0%) had not heard it before. About (95.0%) participants agreed to the fact that it is healthy to comply to a balanced diet plan on regular basis. Most of them (95.0%) knew the value to include whole grain cereals and roughage in their diet chart. About participants (75.0%) were aware of the natural and artificial sources of carbohydrates and (30.%) of them were not aware of their origin. **Discussion:** Although majority of the participants (82.0%) have heard about balanced diet, there are still a few of them who do not follow diet plan themselves frequently at home [5]. According to Juma Al-Kaabi et al (2008), consuming healthy food is a major component for managing all forms of diabetes mellitus, occurring either due to the lack of production insulin resistance, and preventing the serious complication. The aim of dietary modifications is to restore quality of life, caloric management, and preventing long standing hazardous outcomes. [6]. According to a study conducted by Rivellese A A (2007) in Italy concluded that compliance to a comprehensive diet plan is not merely the tuning of dietary components but also the interpretation of the importance of balanced diet. [7] **Conclusion:** Despite basic awareness of the importance of healthy behaviors, the nurses are unable to incorporate their knowledge in the practical and professional lives for their health and wellbeing. Unhealthy outlook of a nurse can be detrimental to the morale, efficiency, adaptability, productivity, and quality of patient care. Nursing instructors can play their crucial role in supervising the adoption of healthy lifestyle patterns so that their subordinates engage in physical exercise, sports, adhere to a balanced diet, combat stress, and improve interpersonal relationships. Furthermore, nursing superintendents can identify and remove

Keywords: metabolic disorders, health-promoting behaviors, balanced diet, physical exercise potential workplace barriers that discourage or prevent nurses from engaging in healthy behaviors.

Corresponding author:**Parveen Akhtar,**

Charge Nurse,

Sheikh Zayed Hospital Rahim Yar Khan,

Email ID: parveentayyab94@gmail.com

QR code



Please cite this article in press Parveen Akhtar et al, *Knowledge And Practices Of Healthy Eating And Sports Among Nurses In Sheikh Zayed Hospital, Rahim Yar Khan., Indo Am. J. P. Sci, 2021; 08(9).*

INTRODUCTION:

The incidence of noncommunicable, metabolic diseases such as hypertension, diabetes, hyperlipidemias has transformed into an epidemic in general population. [1] Nurses are the frontline and most trustable workers to address and respond to this crisis on the behalf of their clients. [2] they hold a special place to counsel and engage their patients in healthy lifestyle behaviors such as adopting a balanced diet, regular physical activity, sleep hygiene, stress management, tobacco and alcohol cessation. Despite the basic awareness nurses are unable to incorporate their knowledge in their practical lives. The following study encompasses all those nurses who are struggling to practice healthy life style patterns despite their hectic job routines.

Health of nurses has been a point of focus in many researches [3]. It has become evident that not only do nurses themselves lag behind in adopting healthy behaviors, but also the environments play an antagonist role. [4] Nurses ought to engage in health promoting activities regarding approach to preventable diseases at individual and collective level under the supervision of their mentors to impregnate health and wellbeing.

While analyzing a demarcation between **health risk behaviors** and **health screening behaviors** must be made [5] Health risk behaviors include smoking,

substance abuse, and alcohol intake. Smoking is as common in nurses as general public. [6] Similarly drug abuse has transformed into a serious issue among nurses in the last few decades. [7]

Health screening behaviors comprises of being physically active and undergoing routine screening for common diseases for timely intervention. [8, 10] Stress and social support can together influence health screening behaviors. Su et al. [9] Health promotion behaviors improves endurance health. [11] These behaviors are critically important in reversing the epidemic of obesity and obesity-related diseases currently afflicting our country.

MATERIALS AND METHODS:

A descriptive study was conducted among 200 enrolled through convenient sampling after seeking informed consent working in different departments of Sheikh Zayed Hospital Rahim Yar Khan. The subjects were both men and women chosen randomly from the different departments of the hospital. A self structured questionnaire consisting of socio-demographic details and knowledge regarding adhering to healthy eating and engaging in sports/physical activity was designed to record the response. A formal approval was taken from the ethical review committee.

Confidentiality and privacy was maintained. Statistical data was analyzed by SPSS 21.

TABLE.1 KNOWLEDGE AND PRACTICES OF HEALTHY DIET AND EXERCISE AMONG PARTICIPANTS (N=200)

Knowledge and Awareness of Balanced Diet	Yes n(%)	No n(%)
Have you ever heard about the term balanced diet?	152 (76.0%)	48(24.0%)
Do you it is important to consume balanced diet daily?	194 (97.0%)	6 (3.0%)
Do you know the portion of fruit/vegetable one should eat everyday?	134(67.0%)	66 (33.0%)
How important fruits and vegetables are in daily diet?	196 (98.0%)	4 (2.0%)
How essential are fats to be consumed daily?	34 (17.0%)	166 (83.0%)
How essential is to consume low fat diet/products?	148 (74.0%)	52 (26.0%)
Do you know the names of types of fats?	140 (70.0%)	60 (30.0%)
Do you prefer lean meat?	130 (65.0%)	70 (35.0%)
Are unseasoned ground nuts good to eat?	172 (86.0%)	28 (14.0%)
Do you know what a starchy food is?	164 (82.0%)	36 (18.0%)
Is it healthy to eat starchy food?	134 (67.0%)	66(33.0%)
Is your diet based on the starchy food item?	68 (34.0%)	132 (66.0%)
Is it essential to add cereals/whole grains/roughage in your diet?	188 (94.0%)	12 (6.0%)
How much sugar should be consumed in a day?	96 (48.0%)	104 (52.0%)
What is the proportion of carbohydrates that make up our daily requirement?	104 (52.0%)	96 (48.0%)
Do you know natural edible sources of carbohydrates?	54 (27.0%)	146(73.0%)
What is DASH diet?	48 (24%)	152 (76%)
How much salt should be taken each day?	124(62%)	76 (38%)
Do you smoke?	72 (36%)	128 (64%)
Do you do brisk walking at least 30 mins/day or 150mins/ week?	64 (32%)	136 (68%)

RESULTS:

The total numbers of participants in the study were 200. The demographic data of the participants shows that among the 200 participants, 88(44.0%) belonged to the age group of 20-30 years, 52(26.0%) between 30-40 years, 24(12.0%) between 40-50 years and 36(18.0%) belong to the age of 50 years and above. More females 148(84.0%) participated than the males 52(16.0%) in this study. The highest number of participants have completed their Diploma 104

(52.0%) followed by Bachelors 76 (38%) and 20 (10%) had Masters Degree.

Table.1 shows the response to the survey questionnaire on awareness of balanced diet. Majority of the participants have heard about balanced diet 152 (76.0%), while 28 (24.0%) have never heard about this term before. About 194 (97.0%) participants think that it is important to consume balanced diet daily.

There were 134 (67.0%) who knew how much portions of fruits and /or vegetables should be taken every day. About 196 (98.0%) participants were aware that it important to eat fruits and vegetables in daily diet. 34 (17.0%) of them responded that fats essential to be included in daily diet. There were 148 (74.0%) participants who are aware that it essential to choose low fat products, and 52 (26.0%) are not aware of it. 140 (70.0%) know about the three kinds of fat. The numbers of participants who opt for lean cut of meats or remove visible fat are 130 (65.0%). A majority of the participants 172 (86.0%) are aware that unsalted nuts and seeds are good to be included in their diet. About 164 (82.0%) knew about starchy foods. Whereas, 134 (67.0%) were not aware of its importance. About 68 (34.0%) of them responded that there is no need to base the main meals around starchy foods. A majority of them 188 (94.0%) were aware that it is essential to include whole grain cereals and pulses in their diet. More than half of the participants 96 (48.0%) are not aware on how much of sugar should be consumed in a day. The participants who knew the daily requirements of carbohydrate were 104 (52.0%). 72 (36%) participants stated that they were actively walking and only 64 (32%) admitted that were walking or performing physical exercise/sports for at least 30 minutes a day or 150 minutes a week.

DISCUSSION:

Although majority of the participants 152 (76.0%) have heard about balanced diet but only 64 (32%) admitted that were walking or exercising /doing sports for at least 30 minutes a day or 150 minutes a week to burn their extra calories. There are still a few of them who have not heard and understood the importance of balanced diet. As per a study conducted in Bijapur, Karnataka(2010), it was seen that the diet was considered to play an important role by 90.68% of the respondents. Khattab MS et al (1999) conducted a study in a Saudi Arabian community, and discovered that only 40% of diabetic patients had good adherence to their diet plan [8]. In the study done by Raj P& Angadi MM (2010), 80% of the respondents reported that they were conscious about their intake but the rest were consuming high carb beverages [9]. The study results are corresponding with Jingran Cao et al, which shows that only 54.10% diabetics were aware that about the diet to be balanced and reasonable [4]. Though 98 % of the participants realised the weightage of including fruits and vegetables in diet, 66 (33.0%) of them were not aware of the portions to be taken daily. 60 (30%) were not aware of the three kinds of fat, 104 (52.0%) did not know the daily requirement of carbohydrates, and 146 (73.0) were not aware of the food sources of carbohydrates. Only 24% knew the abbreviation of DASH diet and 62% knew the daily

salt requirement. These results highlight the fact that there is still a deficiency of knowledge among the nurses regarding nutrition management to prevent both short term and long term complications among diabetic.

Balanced diet and physical activity go hand in hand for promoting health lifestyles and combat metabolic disorders. [12] A sedentary lifestyle increases the risk of mortality and morbidity. [13] Americans spend almost one third of their 24 hours day sitting in front of screens. [14]

The American Heart Association's recommends a minimum of 150 minutes of moderate (or 75 minutes of intense) physical activity per week, in addition to a diet rich in fruits, vegetables, fiber and whole grains, and low in trans fats and sugar. [12] Researches explored non-adherence to these guidelines by the nurses Tucker et al. [15, 16]. Recent literature review has found that the prevalence of obesity in nurses is increasing. [17]

CONCLUSION:

Despite basic awareness of the importance of healthy behaviors, the nurses are unable to incorporate their knowledge in the practical and professional lives for their health and wellbeing. Unhealthy outlook of a nurse can be detrimental to the morale, efficiency, adaptability, productivity, and quality of patient care. Nursing instructors can play their crucial role in supervising the adoption of healthy lifestyle patterns so that their subordinates engage in physical exercise, sports, adhere to a balanced diet, combat stress, and improve interpersonal relationships. Furthermore, nursing superintendents can identify and remove potential workplace barriers that discourage or prevent nurses from engaging in healthy behaviors.

ACKNOWLEDGEMENT

The authors extend their gratitude to the respected colleagues and participants for the valuable time and input.

REFERENCES:

1. Jones DS, Podolsky SH, Greene JA. The burden of disease and the changing task of medicine. *N Engl J Med.* 2012;366(25):2333–2338. [[PubMed](#)] [[Google Scholar](#)]
2. Nurses rank as most honest, ethical profession for 14th straight year. American Nurses Association; [Accessed October 28, 2016]. <http://www.nursingworld.org/2015-NursesRankedMostHonestEthicalProfession>. Published December 21, 2015. [[PubMed](#)] [[Google Scholar](#)]

3. Women's Health Study. [Accessed October 28, 2016];Harvard Medical School, Brigham and Women's Hospital. <http://whs.bwh.harvard.edu/index.html>.
4. Caruso CC. Negative impacts of shiftwork and long work hours. *Rehabil Nurs*. 2014;39(1):16–25. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
5. Committee on Health and Behavior: Research, Practice, and Policy; Board on Neuroscience and Behavioral Health; Institute of Medicine. *Health and Behavior: The Interplay of Biological, Behavioral, and Societal Influences*. Washington, DC: National Academy Press; 2001. [[Google Scholar](#)]
6. Chandrakumar S, Adams J. Attitudes to smoking and smoking cessation among nurses. *Nurs Stand*. 2015;30(9):36–40. [[PubMed](#)] [[Google Scholar](#)]
7. Monroe T, Kenaga H. Don't ask don't tell: substance abuse and addiction among nurses. *J Clin Nurs*. 2011;20(3–4):504–509. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
8. Steptoe A, Wardle J. Health-related behaviour: prevalence and links with disease. In: Kaptein A, Weinman J, editors. *Health Psychology*. Oxford, UK: Blackwell Publishing; 2004. pp. 21–51. [[Google Scholar](#)]
9. Su SY, Chiou ST, Huang N, Huang CM, Chiang JH, Chien LY. Association between Pap smear screening and job stress in Taiwanese nurses. *Eur J Oncol Nurs*. 2016 Feb;20:119–124. [[PubMed](#)] [[Google Scholar](#)]
10. Silva IT, Griep RH, Rotenberg L. Social support and cervical and breast cancer screening practices among nurses. *Rev Lat Am Enfermagem*. 2009;17(4):514–521. [[PubMed](#)] [[Google Scholar](#)]
11. Pender N, Murdaugh C, Parsons MA. *Health Promotion in Nursing Practice*. 6. Upper Saddle River, NJ: Pearson Education, Inc; 2011. [[Google Scholar](#)]
12. The American Heart Association's diet and lifestyle recommendations. American Heart Association; [Accessed October 28, 2016]. http://www.heart.org/HEARTORG/HealthyLiving/HealthyEating/Nutrition/The-American-Heart-Associations-Diet-and-Lifestyle-Recommendations_UCM_305855_Article.jsp. Reviewed August 2015. [[Google Scholar](#)]
13. Rezende LF, Sá TH, Mielke GI, Viscondi JY, Rey-López JP, Garcia LM. All-cause mortality attributable to sitting time: analysis of 54 countries worldwide. *Am J Prev Med*. 2016;51(2):253–263. [[PubMed](#)] [[Google Scholar](#)]
14. Matthews CE, George SM, Moore SC, et al. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr*. 2012;95(2):437–445. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
15. Tucker SJ, Harris MR, Pipe TB, Stevens SR. Nurses' ratings of their health and professional work environments. *AAOHN J*. 2010;58(6):253–267. [[PubMed](#)] [[Google Scholar](#)]
16. Blake H, Malik S, Mo PK, Pisano C. 'Do as I say, but not as I do': are next generation nurses role models for health? *Perspect Public Health*. 2011;131(5):231–239. [[PubMed](#)] [[Google Scholar](#)]
17. Buss J. Associations between obesity and stress and shift work among nurses. *Workplace Health Saf*. 2012;60(10):453–458. 459. [[PubMed](#)] [[Google Scholar](#)]