



CODEN [USA]: IAJ PBB

ISSN : 2349-7750

**INDO AMERICAN JOURNAL OF
PHARMACEUTICAL SCIENCES**

SJIF Impact Factor: 7.187

Available online at: <http://www.iajps.com>

Research Article

**PHARMACOLOGICAL SCREENING AND
PHYTOCHEMICAL EVALUATION OF ANTI-DIABETIC
ACTIVITY OF *TINOSPORA CORDIFOLIA* IN NORMAL AND
STREPTOZOTOCIN INDUCED DIABETIC RATS****M. Pravalika*, P. Geetha Vani, Dr. D. Venkata ramana.**

Department of Pharmacology, Holy Mary Institute of Technology and Science (College of Pharmacy), Keesara - Bogaram - Ghatkesar, Telangana 501301.

Abstract:

*A study of ancient literature indicates that diabetes was fairly well known and well conceived as an entity in India. Plant-based drugs have been used against various diseases since a long time. The nature has provided abundant plant wealth for all the living creatures, which possess medicinal virtues. The essential values of some plants have long been published, but a large number of them have remained unexplored to date. Therefore, there is a necessity to explore their uses and to conduct pharmacognostic and pharmacological studies to ascertain their therapeutic properties. In fact, nowadays, diabetes is a global problem. Hence, the present study aims to open new avenues for the improvement of medicinal uses of *Tinospora cordifolia* (Composite) for the selected area for diabetes. The acute oral toxicity studies of the extracts revealed no toxic effects up to the levels of 2000mg/kg b.wt. The aqueous and alcoholic extracts of 20 and 30mg/kg body weight of *Tinospora cordifolia* was screened for the presence of hypoglycemic and antidiabetic activity. In this study diabetes was induced by a single IP dose Streptozotocin in 72hrs fasted rats. The FBGL was carried on 7th, 14th and 21st day and OGTT was measured on 8th, 15th and 22nd day. Glibenclamide was taken as the standard and the results are quite comparable with it. The studies were indicated that the leaves of *Tinospora cordifolia* are effective in regeneration of insulin secreting β -cells and thus possess antidiabetic activity. The aqueous and alcoholic extracts showed significant effect in decreasing the Fasting blood Glucose level and oral glucose tolerance test of rats and it's also showed good hypoglycemic activity in normal glycemic rats. The preliminary phytochemical analysis of the extracts of *Tinospora cordifolia* revealed the presence of Alkaloids, Flavonoids, Steroids, Tannins, Anthraquinones, Terpenoids and Cardiac glycoside as the possible biologically active principles.*

Keywords: *Tinospora cordifolia*, Streptozotocin, Glibenclamide, FBGL and OGTT.**Corresponding author:****M. Pravalika***Department of Pharmacology,
Holy Mary Institute of Technology and Science (College of Pharmacy),
Keesara - Bogaram - Ghatkesar, Telangana.Email Id-myanapravalika9@gmail.com

QR CODE



Please cite this article in press M. Pravalika et al., *Pharmacological Screening And Phytochemical Evaluation Of Anti-Diabetic Activity Of *Tinospora Cordifolia* In Normal And Streptozotocin Induced Diabetic Rats*, Indo Am. J. P. Sci, 2025; 12(10).

INTRODUCTION:

1.1-Diabetes Mellitus (DM):

Diabetes is one of the most common non-communicable diseases and a serious life-long condition appearing worldwide. The etiology of diabetes is a complex interaction of genetic and environmental factors. It is a heterogeneous group of metabolic disorders characterized physiologically by dysfunction of pancreatic beta cells and deficiency in insulin secretion or insulin activity and clinically by hyperglycemia or impaired glucose tolerance and other manifestable disorders. It is an endocrinological syndrome abnormally having high levels of sugar in the blood. This may be either due to insulin not being produced at all, is not made at sufficient levels, or is not as effective as it should be. Diabetes is still a serious health problem all over the world since it is associated with increased morbidity and mortality rate. When compared with the general population, mortality and morbidity increase in diabetes is mainly due to the associated chronic complications both specific (microvascular) and nonspecific (macrovascular). Since the disease prevails in both genders and in all age groups, the general public has a concern about its control and treatment¹.

1.2-Classification of DM

Diabetes is classified by underlying cause. The most common forms of diabetes are categorized as

Type 1, or insulin-dependent diabetes mellitus (IDDM) - an autoimmune disease in which the body's own immune system attacks the pancreatic beta cells, rendering it unable to produce insulin and **Type 2**, or non-insulin-dependent diabetes mellitus (NIDDM) - in which there is resistance to the effects of insulin or a defect in insulin secretion.

Type 2 diabetes commonly occurs in adults associated with obesity. There are many underlying factors that contribute to the high blood glucose levels in these individuals. An important factor is the resistance to insulin in the body essentially ignoring its insulin secretions. A second factor is the decreased production of insulin by the cells of the pancreas. Therefore, an individual with Type 2 diabetes may have a combination of deficient secretion and deficient action of insulin. In contrast to Type 2 diabetes, Type 1 diabetes most commonly occurs in children and is a result of the body's immune system attacking and destroying the beta cells. The trigger for this autoimmune attack is not clear, but the result is the end of insulin production².

Multiple risk factors for the development of Type 2 diabetes mellitus³:

- Family history (parents with diabetes).
- Obesity (i.e., $\geq 20\%$ over ideal body weight or body mass index $\geq 25\text{kg/m}^2$).
- Habitual physical inactivity.
- Impaired glucose tolerance.

- Hypertension ($\geq 140/90\text{mm Hg}$ in adults).
- High density lipoprotein (HDL) cholesterol $\leq 35\text{mg/dl}$ and/or triglyceride level $\geq 250\text{mg/dl}$.

1.3-History

The term "Diabetes" was first used around 250 B.C. It is a Greek word meaning "to syphon", reflecting how diabetes seemed to rapidly drain fluid from the affected individual. The Greek physician Aretaeus noted that affected individuals passed increasing amounts of urine as if there was "liquefaction of flesh and bones into urine". The complete term "diabetes mellitus" was coined in 1674 by Thomas Willis. Mellitus is Latin for honey, which is how Willis described the urine of diabetics⁵.

Historical accounts reveal that as early as 700-200 BC, diabetes mellitus was a well recognized disease in India and was even distinguished as two types, a genetically based disorder and other one resulting from dietary indiscretion. Ancient Hindu writings document how black ants and flies were attracted to the urine of diabetics. The Indian physician Sushruta in 400 B.C. described the sweet taste of urine from affected individuals, and for many centuries to come, the sweet taste of urine was a key to the diagnosis.

Physicians have observed the effects of diabetes for thousands of years. One of the effects of diabetes is the presence of glucose in the urine (glucosuria). For much of the time, little was known about this fatal disease that caused weight loss of body, extreme thirst, and frequent urination. It was in 1922 that the first patient was successfully treated with insulin. Till the mid-1800s, the treatments offered for diabetes varied tremendously. A breakthrough in the puzzle of diabetes came in 1889. German physicians Joseph von Mering and Oskar Minkowski surgically removed the pancreas from dogs. The dogs immediately developed diabetes. Now that a link was established between the pancreas and diabetes, research focused on isolating the pancreatic extract that could treat diabetes. Dr. Frederick Banting succeeded in his experiments of isolating a pancreatic extract. The diabetic dog was kept alive for eight days by regular injections until supplies of the extract, at that time called "isletin", was exhausted. Experiments on dogs showed that extracts from the pancreas caused a drop in blood sugar, caused glucose in the urine to disappear, and produced a marked improvement in clinical condition.

A young boy, Leonard Thompson, was the first patient to receive insulin treatment in the year 1922 and lived for thirteen years. Over the next 70 years, insulin was further refined and purified. A revolution came with the production of recombinant human

DNA insulin in 1978. Instead of collecting insulin from animals, new human insulin could be synthesized. In 1923, Banting and Macloed were awarded the Nobel Prize for the discovery of insulin. In his Nobel Lecture, Banting concluded the following about their discovery: "Insulin is not a cure for diabetes; it is a treatment."

1.4-Epidemiology⁶

Present status projects that incidence of diabetes is on rise. Present number of diabetics worldwide is 150 million and according to new estimates from researchers at the World Health Organization (WHO), there will be an increase of about 300 million or more by the year 2030 (Warner, 2004). Only in year 2001, about 441,004 deaths were registered and 49,855 of them provoked by diabetes, representing 11.2% of the total population. In United States, diabetes is the sixth leading cause of death. The prevalence of diabetes mellitus is rapidly increasing worldwide and India is estimated to have 31 million diabetics from the total population of the world. Diabetes is predicted to become one of the most common diseases in the world within a couple of decades, affecting at least half a billion people.

The driving force behind the high prevalence of diabetes is the rise of obesity, sedentary lifestyle, consumption of energy rich diet, etc. The diabetes epidemic is accelerating in the developing world, with an increasing proportion of affected people in younger age groups.

The prevalence of Type 2 diabetes is now at epidemic proportions. Type 2 diabetes has a significant impact on the health, quality of life, and life expectancy of patients, as well as on the health care system. Type 2 diabetes accounts for about 90-95 % of population while Type 1 diabetes accounts for about 5 -10% of the total population. In the past, Type 2 was rarely seen in the young, but recent reports describe Type 2 diabetes being diagnosed even in children and adolescent⁷.

6. MATERIALS AND METHODS:

The designing of methodology involves a series of steps taken in a systematic way in order to achieve the set goal(s) under the prescribed guidelines and recommendations. It includes in it all the steps from field trip to the observation including selection and collection of the medicinal plant, selection of dose value, standardization of protocol, usage of instruments, preparation of reagents, selection of specific solvents for extraction, formation of protocols and final execution of the standardized protocol. All this requires good build of mind and a good and soft technical hand to handle the materials and procedure in a true scientific manner.

6.1 Drugs and Chemicals

Drugs and Chemicals used in this study were of analytical grade and of highest purity procured from standard commercial sources in India.

Table No: 6.1 Drugs and Chemicals

S.No	Materials	Company Name
1.	Streptozotocin	Quali Kems Fine Chem Pvt, Ltd, Vadodara.
2.	Methanol	ChangshuYangyuan Chemicals, China.
3.	Alcohol	ChangshuYangyuan Chemicals, China.
4.	Glibenclamide	Sanofi India Ltd, Ankleshwar.

6.2. Instruments

Following instruments were required for the study:

Table No: 6.2- List of Instruments used for study

Name of the instrument	Source
Centrifuge	Dolphin
Digital weighing balance	Horizon
Glucometer	Horizon
Heating mantle	ASGI®
Refrigerator	Videocon
Soxhlet extractor	ASGI®
Condenser	ASGI®
Burette stand	Dolphin
Round bottom flask	ASGI®, Amar
Mixer	Videocon
Oven	ASGI®
Water bath	ASGI®
Stirrer/glass rod	ASGI®
Watch glass	ASGI®

Whatmann filter paper	Manipore microproducts, Ghaizabad.
Butter paper	ASGI®
Spatula	ASGI®
Rubber pipes	ASGI®

6.3. Experimental animals

Healthy adult albino Wistar rats weighing 200-250grams of either sex were selected for the study. Animals were housed in appropriate cages in uniform hygienic conditions and fed with standard pellet diet (Amrul Laboratory Animal Diet) and water ad libitum. They were fasted overnight before the day of experiment, after 72hours of fasting from the day of Streptozotocin introduction. Animals were housed within the departmental animal house and the room temperature was maintained at 27° C. Animal studies had approval of IAEC.

6.4. Plant Material Collection

The leaves of *Tinospora cordifolia* was collected from the local market in Hyderabad and was identified and authenticated from Department of Pharmacognosy. The plant material was cleaned, reduced to small fragments, air dried under shade at room temperature and coarsely powdered in a mixer. The powdered material was stored or taken up for extraction process.⁴⁵

6.5. Preparation of plant extracts:

6.5.1 Preparation of Aqueous Extract:

Dried leaves of *Tinospora cordifolia* were taken about 20gms into 250ml beaker containing 200ml of water. The contents were mixed well and then the mixture was boiled upto 80-90°C for 4-5hrs. Further the extract was filtered with whatmann filter paper. The filtrate was boiled until the concentrated residue is formed. The concentrated product was sealed in sample covers and stored under room temperature and used for further experiment to check the activities.

6.5.2 Preparation of Alcoholic Extract:

Dried leaves of *Tinospora cordifolia* were taken about 20gms into 250ml beaker containing 200ml of Alcohol. The contents were mixed well and then the mixture was boiled up to 50-60°C for 4-5hrs. Further the extract was filtered with whatmann filter paper. The filtrate was boiled until the concentrated residue is formed. The concentrated product was sealed in sample covers and stored under room temperature and used for further experiment to check the activities.⁴⁶

6.6 Preliminary phytochemical analysis of the extracts⁴⁷

The extracts so obtained were subjected to preliminary phytochemical screening. Phytochemical studies were performed to identify the presence of various Phytoconstituents as follows:

6.6.1. Alkaloids

Extracts were dissolved individually in dilute Hydrochloric acid and filtered.

a. Mayer's Test: Filtrates were treated with Mayer's reagent (Potassium Mercuric Iodide). Formation of a yellow coloured precipitate indicates the presence of alkaloids.

b. Wagner's Test: Filtrates were treated with Wagner's reagent (Iodine in Potassium Iodide). Formation of brown/reddish precipitate indicates the presence of alkaloids.

c. Dragendroff's Test: Filtrates were treated with Dragendroff's reagent (solution of Potassium Bismuth Iodide). Formation of red precipitate indicates the presence of alkaloids.

d. Hager's Test: Filtrates were treated with Hager's reagent (saturated picric acid solution). Presence of alkaloids confirmed by the formation of yellow coloured precipitate.

6.6.2. Triterpenoids

a. Salkowski's Test: The extracts were treated with chloroform and filtered separately. The filtrate was treated with few drops of concentrated sulphuric acid, shaken and allowed to stand. If the lower layer turns red, sterols are present. If the lower layer turns golden yellow triterpenes are present.

6.6.3. Saponins

a. Froth Test: The extracts were diluted with distilled water to 20 ml and shaken in a graduated cylinder for 15 mins. The formation of 1 cm layer of foam indicates the presence of saponins.

b. Liberman Burchard Test: The extracts were treated with chloroform and filtered. The filtrates were treated with few drops of acetic anhydride boiled and cooled. Concentrated sulphuric acid was added through the sides of test tube. The formation of brown ring at the junction indicated the presence of steroidal saponins.

6.6.4. Flavonoids

a. Alkaline reagent Test: The extracts were treated with few drops of sodium hydroxide separately. Formation of intense yellow colour less on addition of few drops of dilute acid indicates the presence of flavonoids.

b. Lead acetate Test: The extracts were treated with few drops of lead acetate solution. Formation of yellow precipitate indicates the presence of flavonoids.

6.6.5. Phenolic and Tannins

a. Ferric chloride Test: The extract was treated with few drops of neutral ferric chloride solution. The formation of bluish black colour indicates the presence of phenolics nucleus.

b. Gelatin Test: To the extract, 1%gelatin solution containing sodium chloride was added. The formation of white precipitate indicates the presence of tannins.

c. Vanillin hydrochloride Test: the extracts were treated with few drops of vanillin hydrochloride reagent. The conformation of pinkish red colour indicates the presence of tannins.

6.7. Selection of dose for animal study

The dose considered for the experiment on rats was obtained from conversion of human dose of *Tinospora cordifolia* (3-5 g/kg). The conversion factor of human dose (per 200 g body weight) is 0.018 for rats (Ghosh 1984). Hence the calculated dose for the rats (considering human dose 0.3 and 0.5 g/kg) is 20 and 30 mg/kg. Acute toxicity was done at dose of 2000mg/kg body weight.

6.8. Pharmacological evaluation

Preparation of extracts:

The aqueous and alcoholic extracts of *Tinospora cordifolia* suspended in water in presence of 3%v/v Tween-80 solution.

All the drugs were administered orally for experimental purpose. Each time preparations of the

extracts were prepared when required. The drugs were administered at a constant volume of 10ml/kg for each animal.

6.9. ACUTE ORAL TOXICITY:

The acute oral toxicity of aqueous and alcoholic extracts of *Tinospora cordifolia* was determined by using Albino wistar rats (200-250g) which were maintained under standard conditions. The animals were fasted 12 hour prior to the experiment, up and down procedure OECD guideline no. 425 were adopted for toxicity studies. Animals were administered with single dose of individual extract upto 2000mg/kg and observed for its mortality during 2days and 7days study period (short term) toxicity and observed upto 7days for their mortality, behavioral and neurological profiles.⁴⁸

6.10. Assessment of Anti-diabetic Activity in Normal and Streptozotocin induced Rats

10.1 Assessment of hypoglycemic activity on normal rats.

Table----.Group Classification:

Group	Treatment	Dose(mg/kg)
Group 1	Normal control received distilled water	10ml/kg
Group 2	Standard group received Glibenclamide	10ml/kg
Group 3	Aqueous extract of <i>Tinospora cordifolia</i>	20 mg/kg
Group 4	Aqueous extract of <i>Tinospora cordifolia</i>	30 mg/kg
Group 5	Alcoholic extract of <i>Tinospora cordifolia</i>	20 mg/kg
Group 6	Alcoholic extract of <i>Tinospora cordifolia</i>	30 mg/kg

Procedure:

Animals were divided randomly into six groups of four and each was fasted to overnight. The blood samples were withdrawn by tail vein at 0hour i.e. before I.P administration of extracts/standard/vehicle. Then blood was collected at an interval of 1, 2, 4, and 8 hour after the administration on 0th, 7th, 14th and 21st day respectively according to procedure blood glucose levels were measured by glucometer (ONE TOUCH glucometer).⁴⁹

➤ Oral glucose tolerance test(OGTT) in normal rats:

On the next day (1st, 8th, 15th and 22nd day) after the assessment of hypoglycemic activity OGTT was carried out in same normal animals.

Procedure:

All the animals in each group were administered 2g/kg of glucose one hour after extract/ glibenclamide/ vehicle administration. The blood samples were collected by tail vein at 0 hour, 0.5 hour, 1 hour, 1.5 hour and 2 hour after the administration of glucose load. Blood glucose levels were measured by glucometer on 1st, 8th, 15th and 22nd day respectively.⁵⁰

6.10.2 Assessment of Anti-Diabetic Activity in Streptozotocin Induced Diabetic Rats:

Induction of Diabetes:

Albino wistar rats of either sex weighing 200-250 g were selected for the study. All the animals were allowed free access to water and pellet diet and maintained at room temperature in rat cages.

Streptozotocin was dissolved in normal saline immediately before use. Diabetes was induced in 16 hour fasted rats by single intraperitoneal injection of 120 mg/kg body weight of freshly prepared Streptozotocin in normal saline.

The rats after Streptozotocin were given 5% w/v glucose solution in feeding bottles for next 24 hours in their cages to prevent hypoglycemia. After 72 hours rats with fasting blood glucose levels greater than 200 mg/dl were selected and used for further studies.

All the animals were observed for seven days for consistent hyperglycemia (fasting blood glucose level greater than 200 mg/dl and lesser than 400 mg/dl) and such animals were selected and divided into six groups of four each and used for the study of the following experimental models.⁵¹

Table-----Group Classification:

Group	Treatment	Dose(mg/kg)
Group 1	Normal control received distilled water	10ml/kg
Group 2	Diabetic control received distilled water	10ml/kg
Group 3	Standard group received Glibenclamide	10ml/kg
Group 4	Aqueous extract of <i>Tinospora cordifolia</i>	20mg/kg
Group 5	Aqueous extract of <i>Tinospora cordifolia</i>	30mg/kg
Group 6	Alcoholic extract of <i>Tinospora cordifolia</i>	20 mg/kg
Group 7	Alcoholic extract of <i>Tinospora cordifolia</i>	30 mg/kg

Effect of Aqueous and Alcoholic extracts of *Tinospora cordifolia* on blood glucose levels in Streptozotocin induced diabetic rats:

All the animals of above groups were administered as per treatment protocol mentioned above. The blood samples were collected by retro orbital puncture at 0,1,2,4 and 8 hour after the administration. The treatment was continued for next 22 days. Again blood samples were also collected on 7th, 14th and 21st day after 1 hour administration for sub acute study. Blood glucose level was measured by glucometer at various time intervals.⁵²

Oral glucose tolerance test (OGTT) in Streptozotocin induced diabetic rats:

On the 8th, 15th and 22nd day OGTT was carried out on the same Streptozotocin induced diabetic animals used for assessment of anti-diabetic activity studies.

Procedure:

All the animals in each group were administered 2g/kg of glucose one hour after extract/ Glibenclamide/ vehicle administration. The blood samples were collected by retro orbital puncture at 0 hour, 0.5 hour, 1 hour, 1.5 hour and 2 hour after the administration of the glucose load. The Blood samples were collected by tail vein and its blood

glucose levels were measured by using a glucometer apparatus.⁵³

6.11. Statistical analysis

The values were expressed as mean $\pm$ SEM data was analyzed using one-way ANOVA followed by T-test. Two sets of comparison had made. i.e.

1. Normal control Vs All treated groups.

2. Diabetic Control Vs All treated groups.

Differences between groups were considered significant at $P < 0.001$ and $P < 0.05$ levels.

7. RESULTS:

7.1 Phytochemical screening of *Tinospora cordifolia*.

The present investigation concluded that the isolated compounds from the plant *Tinospora cordifolia* shows the various Pharmacological effects was determined due to the presence of different phytochemical compounds. Further study is needed for the isolation of the constituents present in the plant and its individual pharmacological activity should need to consider and ultimately it should be implemented for the benefit to human beings.

Table 1: Phytochemical screening of *Tinospora cordifolia*.

S.No.	Phytoconstituents	Aqueous	Alcoholic
1.	Alkaloids	+	-
2.	Flavonoids	-	+
3.	Steroids	-	-
4.	Tannins	+	-
5.	Anthraquinones	+	+
6.	Terpenoids	-	-
7.	Cardiac glycoside	+	+

7.2 Acute toxicity testing

Acute toxicity studies revealed that the alcoholic extracts of *Tinospora cordifolia* were safe up to 2000 mg/kg of body weight and approximate LD 50 is more than 2000 mg/kg. No lethality or any toxic reactions was observed up to the end of the study period.

7.3 HYPOGLYCEMIC ACTIVITY IN NORMAL RATS

Fasting Blood Glucose Levels (FBGL) were within the range of 90-105 mg/dl in all the groups at 0day. Repeated treatment with the doses of aqueous and alcoholic extract (100 and 200 mg/kg) significantly decrease the blood glucose level on 7th, 14th and 21st day, indicating that the extract produce significant hypoglycemic activity after repeated administration. Glibenclamide (10mg/kg) also significantly reduced Fasting Blood Glucose Level

(FBGL) after repeated administration as compare to normal control group. Changes in FBGL in different groups after repeated dose administration are summarized in Table No: 10

Repeated administration of both aqueous and alcoholic extracts had significantly ($p < 0.005$) reduced the FBGL on 7th, 15th and 21st day, indicating these extracts can produce hypoglycemia on repeated administration. However hypoglycemic activity was more significant on 7th, 14th and 21st day for Glibenclamide treated as compare with other groups. The results suggest that the both aqueous and alcoholic extracts possess significant hypoglycemic activity after repeated dose administration. The detailed results are summarized in Table No: 10

➤ **Effect of extracts of *Tinospora cordifolia* on fasting blood glucose level (FBGL) in normal rats**

Table No: 10- Effect of extracts of *Tinospora cordifolia* on fasting blood glucose level (FBGL) in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	84.93±3.27	79.28±4.60	74.33±3.11
Glibenclamide	10	79.99±6.10	78.66±2.32	75.89±3.40
AQTC1	20	86.22±3.52	84.24±5.23	80.15±3.75
AQTC2	30	80.36±7.46	78.97±4.64	75.28±1.31
ALTC1	20	70.1±5.86	68.19±3.72	64.78±1.97
ALTC2	30	81.6±5.87	79.20±4.79	76.11±3.62

Values are expressed as mean± S.E.M. n=6. Significant values were compared with $p < 0.005$, normal control Vs all groups. Parent thesis indicates % reduction in BGL.

➤ **Oral glucose tolerance test (OGTT) -**

Both the aqueous and alcoholic extracts of *Tinospora cordifolia* significantly ($P < 0.005$) suppress the rise in FBGL after glucose load (2g/kg) in rats, at first half-an-hour and up to 2hr time period as compare with other groups extract Glibenclamide on 8th, 15th and 22nd day. While aqueous and alcoholic extracts produced significant reduction in FBGL. Glibenclamide (10mg/kg) showed ($P < 0.005$) significant suppression in FBGL rise at first half-an-hour, 1hr and normalized FBGL within 2hr. The detailed results are summarized in Table No: 11

Table No: 11- Effect of extracts of *Tinospora cordifolia* on 8th, 15th and 22nd day in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		8 th day	15 th day	22 st day
Normal control	-	86.15±1.72	90.26±2.84	92.61±1.93
Glibenclamide	10	81.21±1.79	78.27±3.74	75.11±6.50
AQTC1	20	87.26±3.21	84.12±2.36	80.37±6.75
AQTC2	30	79.87±1.33	75.29±1.54	71.47±1.26
ALTC1	20	83.77±0.76	80.35±0.91	75.68±0.76
ALTC2	30	72.18±3.58	69.76±0.44	61.51±0.65

Values are expressed as mean ± S.E.M. n=6. Significant values were compared with $P < 0.005$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

7.4 ANTI-DIABETIC ACTIVITY IN STREPTOZOTOCIN INDUCED DIABETIC RATS

Fasting blood glucose levels (FBGL) in normal rats were in range of 90-100 mg/dl. Treatment with Streptozotocin (120 mg/kg, I.P.) had increased the FBGL to range of 252-266 mg/dl after 72 hours. These values on subsequent days got stabilized by day seven on an average between 255 mg/dl.

Changes in the fasting blood glucose levels in different groups are tabulated in Table No. This data shown that blood glucose level of normal control animals has maintained throughout the study period.

The diabetic control group has shown significant increase in fasting blood glucose levels during this 21st day study period. Glibenclamide (10mg/kg) treated group has shown ($p < 0.05$) significant decrease in fasting blood glucose level during 7th, 14th and 21st day of study period.

➤ **Effect of *Tinospora cordifolia* extracts on antidiabetic activity in Streptozotocin induced diabetic rats**

The animals treated with 100 and 200mg/kg of aqueous and alcoholic of different extracts shown significant decrease ($P<0.05$) in FBGL on 7th, 14th and 21st day of treatment when compare to other groups of animals. The aqueous extracts have reduced more (%) in FBGL when compared to alcoholic extracts except standard group. The detailed results are summarized in Table No: 12

Table No: 12- Effect of extracts of *Tinospora cordifolia* on fasting blood glucose level (FBGL) in Streptozotocin induced diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	81.28±4.69	80.98±7.87	83.74±4.35
Diabetic control	10	298.20±13.21	275.14±10.02	253.31±15.55
Glibenclamide	10	260.11±18.39	249.78±12.05	231.56±15.85
AQTC1	20	370.61±14.71	363.18±19.61	336.87±19.10
AQTC2	30	384.73±12.69	370.88±17.96	353.27±18.33
ALTC1	20	294.78±11.41	276.64±14.88	225.70±15.90
ALTC2	30	210.13±15.64	191.21±16.86	165.90±10.38

Values are expressed as mean $\pm$ S.E.M. n=6. Significant values were compared with $P<0.05$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

➤ **Oral glucose tolerance test (OGTT) on 8th, 15th and 22nd day-**

Both the aqueous and alcoholic extracts of *Tinospora cordifolia* are significantly ($P<0.05$) suppress the rise in FBGL after glucose load (2g/kg) in rats, at first half-an-hour and up to 2hr time period as compare with other groups extract Glibenclamide on 8th, 15th and 22nd day. While aqueous and alcoholic extracts produced significant reduction in FBGL. Glibenclamide (10mg/kg) showed ($P<0.05$) significant suppression in FBGL rise at first half-an-hour, 1hr and normalized FBGL within 2hr. The detailed results are summarized in Table No: 13.

Table No: 13- Effect of extracts of *Tinospora cordifolia* on 8th, 15th and 22nd day in Diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level (mg/dl)		
		8 th day	15 th day	22 st day
Normal control	-	87.75±1.63	84.59±2.87	90.99±1.58
Diabetic control	10	288.10±12.69	297.00±19.55	326.57±14.99
Glibenclamide	10	368.78±18.10	310.88±11.73	290.41±18.55
AQTC1	20	260.36±16.62	243.70±16.40	218.36±11.90
AQTC2	30	290.84±11.30	286.78±13.78	275.98±19.75
ALTC1	20	266.42±18.11	223.56±17.87	198.63±17.99
ALTC2	30	356.85±12.61	335.76±15.28	290.66±10.86

Values are expressed as mean $\pm$ S.E.M. n=6. Significant values were compared with $P<0.05$. Normal control Vs all groups. Paranthesis indicates % reduction in BGL.

8. DISCUSSION:

Despite the fact that diabetes has high prevalence, morbidity and mortality globally, it is regarded as non curable but controllable disease. Different synthetic drugs, plant remedies and dietary modification play an effective role in the reduction of the suffering that it causes. The potential role of medicinal plants as antidiabetic agents has been reviewed by several authors. In order to identify the plants with antidiabetic properties various plants have been tested *in-vivo* using animal models, for example rats, against the complications caused by inducers of diabetes, and it has been established that many plants possesses the potential to lower the fasting blood glucose levels and besides help in improving other diabetic complications. The

sustained reduction in hyperglycemia automatically decreases the risk of other major complications of diabetes. Effective glucose control is the key for preventing or reversing the diabetic complications and improving the quality of life of the diabetics.

Many natural active compounds have been isolated from plants of different species. These active principles are complex Alkaloids, Flavonoids, Steroids, Tannins, Anthraquinones, Terpenoids, Cardiac glycoside and others. These compounds have been shown to produce potent hypoglycemic, anti-hyperglycemic and glucose suppressive activities. These effects might be achieved by facilitating insulin release from pancreatic β -cells, inhibiting glucose absorption in gut, stimulating glycogenesis in liver and/ or increasing glucose

utilization by the body. These compounds may also exhibit Anti-Inflammatory, Antibacterial, Antifungal and Cardio protective activities, and restore enzymatic functions, repair and regeneration of pancreatic islets and the alleviation of liver and renal damage.

Crude aqueous and alcoholic extracts of leaves of *Tinospora cordifolia* at a dose of 20 and 30mg/kg showed significant effect on the glucose tolerance of rats and it also showed reduction in the fasting blood glucose levels of the normoglycaemic rats, thus revealing the hypoglycemic nature of the extracts. The effect was more pronounced for both extracts. These findings indicate that the extracts might be producing hypoglycemic effect by a mechanism independent from the insulin secretion e.g. by the inhibition of endogenous glucose production or by the inhibition of intestinal glucose absorption.

Streptozotocin is one of the chemical agents used to induce diabetes mellitus in animals. It induces diabetes by dose dependent destruction of β -cells of islets of langerhans. It is a generator of free radicals of oxygen which cause extensive DNA damage. It was observed that single intravenous dose of Streptozotocin exhibited significant hyperglycemia. Excessive hepatic glycogenolysis and gluconeogenesis associated with decreased utilization of glucose by tissues is the fundamental mechanism underlying hyperglycemia in the diabetic state. As the hyperglycemia induced by Streptozotocin falls under category of mild diabetes and may reverse after a few weeks, the hypoglycemic effect of the plant in hyperglycemic rats was studied during 22 days treatment. The difference observed between the initial and final fasting serum glucose levels of extract treated hyperglycemic rat's revealed antihyperglycemic effect of leaves of *Tinospora cordifolia* throughout the period of study. The effect of the extracts was compared to that of reference standard, Glibenclamide and was found to be significant.

Phytochemical analysis of extracts of leaves of *Tinospora cordifolia* revealed the presence of secondary metabolites that have been shown to possess antidiabetic effect in other plants. Flavonoids, alkaloids and Steroids which were responsible for the antidiabetic effect in other plants were also detected in the extracts of this plant. The presence of phenols in the plant could also be responsible for the antidiabetic effect have been shown to prevent the destruction of β -cells by inhibiting the peroxidation chain reaction and thus they may provide protection against the development of diabetes. Extracts of leaves of *Tinospora cordifolia* appear to be attractive materials for further studies leading to possible drug development for diabetes. Development of phytomedicine is relatively inexpensive and less time consuming; it is more suited to our economic

conditions than allopathic drug development which is more expensive and spread over several years.

9. SUMMARY:

1. The leaves of *Tinospora cordifolia* for this project work were procured locally.
2. The leaves of *Tinospora cordifolia* were successively extracted with water and alcohol.
3. Therapeutic dose of the extracts were calculated after carrying acute oral toxicity studies in rats.
4. Extracts were tested for their anti-diabetic activity in normal and Streptozotocin induced diabetic rats.
5. The following parameters were assessed:
 - fasting blood glucose levels
At 7th, 14th and 21st day in normal and Streptozotocin induced rats.
 - Oral Glucose Tolerance Test
At 8th, 15th and 22nd day in normal and Streptozotocin induced rats.
6. Aqueous and Alcoholic extracts of *Tinospora cordifolia* (20 and 30 mg/kg) showed significant effect in blood glucose lowering activity and improved oral glucose tolerance test (OGTT) in short term (7th day) and long term (14th and 21st day) repeated administration in normal and Streptozotocin induced diabetic rats.
7. The above studies showed that Aqueous and Alcoholic extracts of *Tinospora cordifolia* had potent anti-diabetic activity on repeated administration.

10. CONCLUSION:

The study was performed to find out the beneficial effects of two different extracts of leaves of *Tinospora cordifolia* in normoglycaemic rats and Streptozotocin induced diabetic rats and the results reveal that the plant has beneficial effects on blood glucose levels.

In current scenario, herbs are the potent sources of medicines used in the treatment of various disease and disorders. Since, plants are used as medicine there is prompt need of evaluation of plant species, therefore, the present work was conceived to evaluate the phytochemical and pharmacological screening of leaves of *Tinospora cordifolia*. The Phytochemical evaluation has revealed the presence of Alkaloids, Flavonoids, Steroids, Tannins, Anthraquinones, Terpenoids, Cardiac glycoside. The aqueous and alcoholic extracts had hypoglycemic activity because the presence of flavonoids which are rich in treatment of hypoglycemia with less side effects. Flavonoids might be producing hypoglycemic effect by a mechanism independent from insulin secretion e.g. by the inhibition of endogenous glucose production or by the inhibition of intestinal glucose absorption.

The present study *Tinospora cordifolia* of both aqueous and alcoholic extracts was showed significant effect on glucose tolerance and also showed reduction in fasting blood glucose levels in normal diabetic rats.

The data of the blood glucose level of rats treated with Streptozotocin (150mg/kg body weight) produced diabetes within 72 hours. After 72 hours of Streptozotocin administered the blood glucose levels of rats were observed. It was observed that significant lowering of sugar in aqueous and alcoholic extract. The administration of different extracts at a dose of 20 and 30 mg/kg showed significant anti-hyperglycemic effect at 22nd day which was evident from the 7th day onwards as compared to standard. The aqueous and alcoholic extract of *Tinospora cordifolia* has showed better anti-hyperglycemic effect of the extract on the fasting blood sugar levels on diabetic rats are shown in table. The decreasing blood glucose levels are comparable with that of 10 mg/kg of Glibenclamide. The Glibenclamide (10 mg/kg body weight) shows significant effect on compare to the initial and more significant effect on the 22nd Day compare to the initial. The aqueous and alcoholic extracts of 20 and 30mg/kg body weight shows significant ($P < 0.05$), effect.

Results of anti-diabetic activity in normal and Streptozotocin induced rats the extracts established the scientific basis for the utility of these plants in the treatment of diabetes. The extracts have shown significant reduction in blood glucose levels in normal and Streptozotocin induced diabetic rats and produced maximum anti-diabetic activity and is higher than the hypoglycaemic activity of Glibenclamide in the diabetic rats. In glucose loaded animals, the drug has reduced the blood glucose to the normal levels. It is possible that the drug may be acting by potentiating the pancreatic secretion or increasing the glucose uptake. In conclusion, these extract showed significant anti-diabetic effect in normal and diabetic rats after administration. Thus the claim made by the traditional Indian systems of medicine regarding the use of these plants in the treatment of diabetes stands confirms.

ACKNOWLEDGEMENT

The Authors are thankful to the Management and Principal, Holy Mary Institute of Technology and Science (College of Pharmacy), Keesara - Bogaram - Ghatkesar, Telangana, Telangana, for extending support to carry out the research work. Finally, the authors express their gratitude to the Sura Pharma Labs, Dilsukhnagar, Hyderabad, for providing research equipment and facilities

11. REFERENCES:

1. Consultation, W. H. O. Definition, diagnosis and classification of diabetes mellitus and its complications. Vol. 1. Part, 1999.

2. Alberti, Kurt George Matthew Mayer, and PZ ft Zimmet. "Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation." *Diabetic medicine* 15.7 (1998): 539-553.
3. Gress, Todd W., et al. "Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus." *New England Journal of Medicine* 342.13 (2000): 905-912.
4. Ahmed, Awad M. "History of diabetes mellitus." *Saudi medical journal* 23.4 (2002): 373-378.
5. Zimmet, Paul Z., Daniel J. McCarty, and Maximilian P. de Courten. "The global epidemiology of non-insulin-dependent diabetes mellitus and the metabolic syndrome." *Journal of Diabetes and its Complications* 11.2 (1997): 60-68.
6. American Diabetes Association. "Type 2 diabetes in children and adolescents." *Pediatrics* 105.3 (2000): 671-680.
7. Saltiel, Alan R., and C. Ronald Kahn. "Insulin signalling and the regulation of glucose and lipid metabolism." *Nature* 414.6865 (2001): 799-806.
8. Browning, Jeffrey D., and Jay D. Horton. "Molecular mediators of hepatic steatosis and liver injury." *Journal of Clinical Investigation* 114.2 (2004): 147-152.
9. Aronoff, Stephen L., et al. "Glucose metabolism and regulation: beyond insulin and glucagon." *Diabetes Spectrum* 17.3 (2004): 183-190.
10. Ross, E. J., and D. C. Linch. "Cushing's syndrome—killing disease: discriminatory value of signs and symptoms aiding early diagnosis." *The Lancet* 320.8299 (1982): 646-649.
11. Levin, Marvin E., Vincenza C. Boisseau, and Louis V. Avioli. "Effects of diabetes mellitus on bone mass in juvenile and adult-onset diabetes." *New England Journal of Medicine* 294.5 (1976): 241-245.
12. Sherwin, Robert, and Philip Felig. "Pathophysiology of diabetes mellitus." *The Medical clinics of North America* 62.4 (1978): 695-711.
13. Faber, O. K., and C. Binder. "C-peptide response to glucagon: a test for the residual β -cell function in diabetes mellitus." *Diabetes* 26.7 (1977): 605-610.
14. Ioannidis, Ioannis. "Pathophysiology of Type 1 diabetes." *Diabetes in Clinical Practice: Questions and Answers from Case Studies* 31 (2007): 23.
15. Scheen, A. J. "Pathophysiology of type 2 diabetes." *Acta Clinica Belgica* 58.6 (2003): 335-341.

16. Vambergue, A., et al. "[Pathophysiology of gestational diabetes]." *Journal de gynecologie, obstetrique et biologie de la reproduction* 31.6 Suppl (2002): 4S3-4S10.
17. Inzucchi, Silvio E. "Diagnosis of diabetes." *New England Journal of Medicine* 367.6 (2012): 542-550.
18. Turner, R. C., et al. "Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33)." *lancet* 352.9131 (1998): 837-853.
19. Brownlee, M., and A. Cerami. "The biochemistry of the complications of diabetes mellitus." *Annual review of biochemistry* 50.1 (1981): 385-432.
20. Pari, L., and G. Saravanan. "Antidiabetic effect of Cogent db, a herbal drug in alloxan-induced diabetes mellitus." *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 131.1 (2002): 19-25.
21. Day, Caroline. "Traditional plant treatments for diabetes mellitus: pharmaceutical foods." *British Journal of Nutrition* 80.01 (1998): 5-6.
22. Smith, George Davey, et al. "Genetic epidemiology and public health: hope, hype, and future prospects." *The Lancet* 366.9495 (2005): 1484-1498.
23. Shahida Akter, Hanif Ali, Ali A. Shati, Mohammad Y. Alfaifi, Serag Eldin I. Elbehairi, R. Z. Sayyed, Tanzima Yeasmin. Antidiabetic activity of methanolic extract of *Hibiscus sabdariffa* Linn. fruit in alloxan-induced Swiss albino diabetic mice. *Open Agriculture* 2024; 9: 20220243.
24. Kolajo Adedamola Akinyede, Habeebat Adekilekun Oyewusi, Gail Denise Hughes, Okobi Eko Ekpo, and Oluwafemi Omoniyi Oguntibeju. In Vitro Evaluation of the Anti-Diabetic Potential of Aqueous Acetone *Helichrysum petiolare* Extract (AAHPE) with Molecular Docking Relevance in Diabetes Mellitus. *Molecules* 2022, 27, 155.
25. Nithiyaa Perumal, Meenakshii Nallappan, Shamarina Shohaimi, Nur Kartinee Kassim, Thiam Tsui Tee, Yew Hoong Cheah. Synergistic antidiabetic activity of *Taraxacum officinale* (L.) Weber ex F.H. Wigg and *Momordica charantia* L. polyherbal combination. *Biomedicine & Pharmacotherapy* 145 (2022) 112401.
26. Yuting Zhu , Hongwei Gao, Shanhao Han, Jianhui Li, Qiqin Wen, Bo Dong. Antidiabetic activity and metabolite profiles of ascidian *Halocynthia roretzi*. *Journal of Functional Foods* 93 (2022) 105095.
27. Aligwekwe A. Ugochukwu and Idaguko C. Anna, Phytochemical Evaluation and Anti-Diabetic Effects of Ethanolic Leaf Extract of *Petersianthus macrocarpus* on Streptozotocin-Induced Diabetic Rats. *JAMMR*, 33(3): 39-47, 2021.
28. Lia Ardiana, Rani Sauriasari, Berna Elya. Antidiabetic Activity Studies of White Tea (*Camellia sinensis* (L.) O. Kuntze) Ethanolic Extracts in Streptozotocin-nicotinamide Induced Diabetic Rats. *Pharmacogn J.* 2018; 10(1):186-189.
29. Hanae Naceiri Mrabti, Nidal Jaradat, Ismail Fichtali, Wessal Ouedrhiri, Shehdeh Jodeh, Samar Ayeshe, Yahia Cherrah and My El Abbes Faouzi. Separation, Identification, and Antidiabetic Activity of Catechin Isolated from *Arbutus unedo* L. Plant Roots. 12 April 2018.
30. Ram Niwas Jangir, Gyan Chand Jain. Evaluation of Antidiabetic Activity of Hydroalcoholic Extract of *Cassia fistula* Linn. pod in Streptozotocin-Induced Diabetic Rats. *Pharmacogn J.* 2017; 9(5): 599-606.
31. Ameebahen B. Patel , Dharmeshkumar D. Prajapati, Yogesh Patel. Antidiabetic activity of *Moringa oleifera* Lam. Algerian Journal of Natural Products 5:2 (2017) 446-453.
32. Md. Uzzal Haque, Samiron Sana. Antidiabetic activity of plants, fruits and vegetables: a review. *Int J Biol Med Res.* 2016;7(1):5452-5458.
33. Juan Wang, Lirong Teng, Yange Liu, Wenji Hu, Wenqi Chen, Xi Hu, Yingwu Wang, and Di Wang. Studies on the Antidiabetic and Antinephritic Activities of *Paecilomyces hepialid* Water Extract in Diet-Streptozotocin-Induced Diabetic Sprague Dawley Rats. 1 February 2016.
34. Dr. D. Radhika and R. Priya. Assessment Of Anti-Diabetic Activity Of Some Selected Seaweeds. *ejbps*, 2015, Volume 2, Issue 6, 151-154.
35. Dyah Ratna Wulan, Edi Priyo Utomo, and Chanif Mahdi. Antidiabetic Activity of *Ruellia tuberosa* L., Role of α -Amylase Inhibitor: In Silico, In Vitro, and In Vivo Approaches. 28 September 2015.
36. Khalid Ghazanfar, Bashir A. Ganai, Seema Akbar, Khan Mubashir, Showkat Ahmad Dar, Mohammad Younis Dar, and Mudasar A. Tantry. Antidiabetic Activity of *Artemisia amygdalina* Decne in Streptozotocin Induced Diabetic Rats. 21 May 2014.
37. Abdul Vahab A and Jyoti Harindran. Anti-diabetic Activity of *Terminalia catappa* Bark Extracts in Alloxan Induced Diabetic Rats. 16/05/2014.
38. Taofik O. Sunmonu and Anthony J. Afolayan. Evaluation of Antidiabetic Activity and Associated Toxicity of *Artemisia afra* Aqueous Extract in Wistar Rats. 20 May 2013.
39. Anil Kamboj, Sunil Kumar and Vipin Kumar. Evaluation of Antidiabetic Activity of

- Hydroalcoholic Extract of *Cestrum nocturnum* Leaves in Streptozotocin-Induced Diabetic Rats. 18 August 2013.
40. Jude E Okokon, Bassey S Antia and John A Udobang. Antidiabetic activities of ethanolic extract and fraction of *Anthocleista djalensis*. Asian Pac J Trop Biomed. 2012 Jun; 2(6): 461–464.
41. Himanshu Misra, Manish Soni, Narendra Silawat, Darshana Mehta, B. K. Mehta, and D. C. Jain. Antidiabetic activity of medium-polar extract from the leaves of *Stevia rebaudiana* Bert. (Bertoni) on alloxan-induced diabetic rats. J Pharm Bioallied Sci. 2011 Apr-Jun; 3(2): 242–248.
42. Mukesh S. Sikarwar and M. B. Patil. Antidiabetic activity of *Crateva nurvala* stem bark extracts in alloxan-induced diabetic rats. J Pharm Bioallied Sci. 2010 Jan-Mar; 2(1): 18–21.
43. Fenglin Li, Qingwang Li, corresponding Dawei Gao, and Yong Peng. The Optimal Extraction Parameters and Anti-Diabetic Activity of Flavonoids from *Ipomoea Batatas* Leaf. Afr J Tradit Complement Altern Med. 2009; 6(2): 195–202.
44. https://en.wikipedia.org/wiki/Tinospora_cordifolia
45. Gobinath R, Parasuraman S, Sreeramanan S, Chinni SV. Antidiabetic and antihyperlipidemic activities of methanolic extract of leaves of *Coccinia grandis* in diabetic rats. Int J Res Pharm Sci. (2020) 11:2324–31. doi: 10.26452/ijrps.v11iSPL4.4460.